

# FLEA BEETLE DAMAGE TO GARDEN CROPS

LEONARD HASEMAN

Most gardeners fail to recognize the destructive work of small flea beetles. This is due to the small size of the pest and its ability to quickly leave the scene of action when disturbed. The all too common injury in the form of holes the size of pin pricks on such crops as potato, tomato, radish, bean, eggplant, mustard, cabbage, and other crops, is the work of flea beetles.

There are several different species of flea beetles which attack the different garden crops. They vary in color, some being black, some greyish, some brown, and some having dark and light stripes. And they may vary in size, but typical ones are small, about the size of a pinhead, and equipped with jumping hind legs, from which they get the name flea beetle.

Some flea beetles are rather general feeders, attacking several different crops while others may be restricted primarily to a single crop. In some cases, the adult beetles attack the foliage of the crop, while the small, white grub bores in the stems or feeds on the roots below ground. In most cases, the winter is spent in the adult stage hidden away under leaves, grass, or other protection in the garden along fences or in adjoining waste areas. When conditions are favorable for the pest, it may move into a garden from waste ground in such numbers as to seriously damage or destroy crops. Generally speaking, the most serious damage is done early in the season, due to the heavy feeding of the beetles before they mate and lay their eggs. The life history of the different species varies, some having only one generation a year while others may develop more rapidly.

**Control.**—With pests of this type, which also may feed and breed on various weeds botanically closely related to the garden crops, it is important that the grower keep down weeds in and near the garden plot. Also, if a new tract of ground is to be used for a garden it should be plowed in late summer or fall prior to spring planting to eliminate weeds and harbors of the flea beetles. In spring, prior to planting the different crops, cultivate the ground to keep down weeds which may attract the beetles to the garden.

When heavy infestations appear, make prompt applications of DDT, methoxychlor, pyrethrum, rotenone or derris. As soon as injury appears, make application and repeat in a week if the pest continues abundant.

Some growers prefer to make dust applications. In such cases use 3 per cent DDT or methoxychlor powder. If applied when the plants are moist with dew, dust applications stick to the foliage better. Apply the spray or dust so that it coats both the lower side and the top of the leaves. Repeat dust applications several times at intervals of a week or less.

In case of mustard and other leafy vegetables, use a non-poisonous insecticide containing pyrethrum or derris. To protect cabbage, tomato, tobacco, and other plants in the seedbed, keep the bed covered with cheesecloth. When such plants are being set out, dip the tops in a solution containing one of the above insecticides to protect them until the plants get a good start. The secret of success in fighting flea beetles is to be ready to spray or dust when they first show up.

While it is important that gardeners keep everlastingly on the lookout for outbreaks of these and other crop pests and be ready to use sprays or dusts promptly, they must not overlook the far-reaching importance of garden sanitation and proper cultural practices. While the above insecticides are very effective, cultural practices or other preventive measures which will reduce garden insect numbers should be used. Prompt disposal of crop residues, with the insects still attacking them, will help. Fall or winter cleanup and spading or plowing of the garden will often go far in reducing next year's insect losses. Keeping down weeds around the garden and clean cultivation to get rid of weeds in the garden will serve the same end as the use of sprays and dusts.