

# **Peripartum Cardiomyopathy**

## **Background**

### 1. Definition

- Heart failure without identifiable cause, with onset in late pregnancy or early postpartum

## **Pathophysiology**

### 1. Pathology of disease

- Unclear
- Suspected factors include myocarditis (viral), genetic, increased pregnancy-related hemodynamic changes

### 2. Incidence, prevalence

- Varies worldwide
- 1:15,000 in US
- Higher in other areas, especially Haiti & some African nations (as high as 1:100 in some regions)

### 3. Risk factors

- Age
  - Higher risk with age >30 years
- Genetic
  - African-Americans at higher risk
  - Some increased risk within families with history of peripartum cardiomyopathy - genetic vs. exposure?
- Pregnancy-related factors
  - Multiparity
  - Multiple gestation
  - History of pregnancy-induced hypertension, preeclampsia, or eclampsia
  - Long term use of beta-blockers
- Environmental
  - Cocaine use during pregnancy

### 4. Morbidity / mortality

- Mortality
  - 6-23% in US based case series
- Cardiac transplant rate
  - 4-7% eventually require heart transplant
- One-half of patients reach EF >50% by 2 years
  - Most of this recovery occurs in first 6 mos
- Potential risk in future pregnancies
  - Further pregnancies may worsen EF - risk may exist even if cardiac function had normalized

## **Diagnostics**

### 1. History

- Dyspnea, cough, orthopnea, PND, hemoptysis, fatigue, occasionally abdominal pain, palpitations

### 2. Physical exam

- Displaced PMI, gallop, edema

### 3. Diagnostic testing

- Echo and EKG most important
- Laboratory evaluation
  - Enzymes- usually normal
  - Consider viral cultures/titers (including coxsackie B)
- Diagnostic imaging
  - Echocardiogram
  - LV enlargement
  - Contractility impaired
  - Measurements of ventricle size, EF important
- Consider CXR if post-delivery
- Other studies
  - EKG
    - May show LVH, sinus tachycardia, nonspecific ST changes
    - Occasionally other arrhythmia, Q waves
  - Less commonly used, may be needed for severe case:
    - Heart Catheterization
    - Endomyocardial biopsy
  - Tests needed to exclude other diagnostic concerns:
    - Spiral CT or VQ scan to rule-out PE

### 4. Criteria for diagnosis

- Time of onset
  - After 36 weeks of pregnancy or first 5 mos. postpartum
  - Those with idiopathic cardiomyopathy before 36 wks excluded from this definition, but may have similar prognosis & course
- No identifiable cause
- No signs/symptoms of heart disease prior to last month of pregnancy
- Signs of left ventricular systolic dysfunction (LSVD) on echo (EF<45%)

## Differential Diagnosis

### 1. Key DDx

- Pulmonary embolus
- Pneumonia
- CHF from other causes
- Preexisting cardiac disease

### 2. Extensive DDx

- Pulmonary edema related to preeclampsia or eclampsia or treatment for these conditions

## Therapeutics

### 1. Acute treatment

- Similar to other heart failure
  - Oxygen
  - Diuretics
  - Na restriction
  - Digoxin
  - Beta blockers
  - ACE-I or ARB if post-delivery

- Anticoagulation (esp. for severe dilation)
  - Avoid warfarin prior to delivery
2. Further management (24 hrs)
    - Cardiac monitoring
    - Treat arrhythmias
    - Pulmonary emboli more common
  3. Long-term care
    - Monitor with follow-up echos (after 2-3 wks, again q 6 mos. until complete recovery or plateau in improvement)
    - Severely decreased EF without improvement may require heart transplant
    - Document normalized LV function prior to attempting future pregnancy & counsel about potential risks
    - Counsel those with persistent LVSD to avoid future pregnancy

### **Follow-Up**

1. Admit to hospital
  - All patients with new onset heart failure or decompensation
2. Refer to specialist

### **Prognosis**

1. Most who regain EF >50%, do so within 6 mos
2. Best prognosis for regaining EF is in those with initial EF >30%

### **References**

1. Jacobs AG, Bales AC, Lang RM. "Peripartum Cardiomyopathy" in UpToDate, Rose BD (Ed), UpToDate, Waltham, MA, 2007.
2. Sliwa K, Fett J, Elkayam U. "Peripartum Cardiomyopathy" in Lancet 2006; 368: 687-693.
3. Elkayam U, Akhter MW, Singh H, Khan S, Bitar F, Hameed A, Shotan A. "Pregnancy-Associated Cardiomyopathy: Clinical Characteristics and a Comparison Between Early and Late Presentation" in Circulation 2005; 111: 2050-2055.
4. Elkayam U, Tummala P, Rao K, Akhter M, Karaalp I, Wani O, Hameed A, Gviazda I, Shotan A. "Maternal and Fetal Outcomes of Subsequent Pregnancies in Women with Peripartum Cardiomyopathy" in NEJM 2001; 344: 1567-1571.

**Authors: Robin DeMuth, MD, & Robert Darios, MD, Michigan State University-Sparrow Hospital FPRP**

**Editor: David Wakulchik, MD, Aultman FMRP, OH**