378.7M71
X645

UMLD

UM Libraries Depository

104034103006

Gladney.
Foundation of Geometry.

This Thesis Has Been

MICROFILMED

Negative No. T-

542

Form 26

The Foundations of Geometry
by
A. G. Hadwiger

The Foundations of Geometry.
Geometry has been called
the science of in-direct
measurement, and as such
is founded on certain def-
initions, postulates, and
some assumptions or axioms
which are said to be self-
evident. It is a physical
science idealized. The old-
est and best known form
of the science is the
Euclidean Geometry which
was developed by the Greeks.
In this Geometry we
find the following def-
initions of terms, postu-

lates, and axioms.

Definitions

- (1) A point is that which has no magnitude.
- (2) A line is that which has length but no breadth, or thickness.
- (3) The ends of lines are points.
- (4) The right line is such that any part will, however placed, lie wholly on any other part, if its extremities are made to fall on that other part.
- (5) A surface is that which has length and breadth.

without thickness.

- (6) The extremities of a surface are lines.
- (7) A plane surface is a surface in which any two points being taken the straight line that joins them lies wholly in the surface.
- (8) A plane angle is the difference in direction of two straight lines which intersect and lie in the same plane.
- (9) When the lines which form a plane angle are right lines, the angle is called rectilinear.

- (10) When two right lines intersect in a plane making the two adjacent angles equal, each of those angles is called a right-angle, and the lines are said to be perpendicular.
- (11) An obtuse angle is an angle which is greater than a right angle.
- (12) An acute angle is an angle which is less than a right angle.
- (13) A limit is the extremity of anything.
- 14 A figure is that which is surrounded by one or by several

Limits.

- (15) A circle is a plane figure contained by one line which is called the circumference and is such that all right lines drawn from a certain point within the circle to the circumference are equal to each other.
- (16) This point in the circle is called the centre.
- 17 A diameter of a circle is a right line drawn through the centre and terminated both ways by the circumference.
- 18 A semi-circle is the figure made by a diameter and

one of the equal parts of the circumference made by the diameter.

(19) A segment of a circle is the figure contained by a right line and either one of the two parts of the circumference made by the right line.

(20) A figure made by right lines is a rectilinear figure.

(21) If the figure is made by three lines, it is a triangle.

(22) If the figure is made by four lines it is a quadrilateral.

(23) A figure which is enclosed by more than four lines is a polygon.

24 An equilateral triangle is a triangle which has three equal sides.

(25) An isosceles triangle has two equal sides.

26 A scalene triangle is one which has three unequal sides.

27 A triangle which has one right angle is a right triangle.

28 A triangle which has three acute angles is an acute angled triangle.

29 A Triangle which has one obtuse angle is an obtuse angled triangle.

30 A square is a quadrilateral with four equal sides and all of its angles right angles.

(31) A rectangle is a quadrilateral with all of its angles right angles but not all of its sides equal.

(32) A rhombus is a quadrilateral with four equal sides but with its angles not right angles.

33 A parallelogram is a quadrilateral whose opposite sides are equal, and whose oppo-

side angles are equal, but
with its angles not right
angles.

(34) All other four sided figures
are called quadrilaterals.

Postulates.

Let it be granted that:

(1) A straight line can be
drawn from any point in
space to any other point
in space.

(2) Any finite straight line
can be produced indefin-
itely.

(3) A circle may be described
about any point as a centre

and have a radius equal to any finite straight line.

Axioms

- (1) Magnitudes that are equal to the same magnitude are equal to each other.
- (2) If to equal magnitudes equal magnitudes are added the sums are equal magnitudes.
- (3) If from equal magnitudes we take equal magnitudes the remainders will be equal magnitudes.
- (4) If to unequal magnitudes we add equal magnitudes

These are unequal magnitudes

(5) If from unequal magnitudes we take equal magnitudes the remainders are unequal magnitudes.

(6) Magnitudes which are double the same magnitudes are equal.

(7) Magnitudes which are halves of the same magnitudes are equal.

(8) Magnitudes that can be made to coincide are equal.

(9) The whole is greater than any of its parts.

(10) All right angles are equal.

(11) If two right lines are in-

intersected by a third right line which forms with them two interior angles upon the same side of this line, and if the sum of these two interior angles be less than two right angles, these two lines, if prolonged indefinitely, and by intersecting on that side of the line where the sum of the two interior angles is less than two right angles.

"2) Two straight lines can not enclose a space.

These are the definitions, postulates, and axioms,

181

upon which the Euclidean
Geometry is founded.

We call the Euclidean Geometry
on account of its origin and
to distinguish it from other
Geometries called non-Euclidean
and projective Geometry. These
latter Geometries have been
developed in modern times
and are not founded on
the same axioms.

By Geometry alone we mean
to include both the Euclidean
and the non-Euclidean Geometries.

And we shall find that Geo-
metry is founded upon two
kinds of axioms. (1) Those axioms

without which experience of the
 the subject matter would
 be impossible. '2) These axioms
 which depend upon experience.
 The first kind we call
 a priori axioms the second
 kind we call empirical axioms.

Now of the first
 seven and ninth Euclidean axioms
 we say that they are of the
 first or a priori kind. For
 they are necessary to the existence
 of mathematics. Consider the
 first which can be stated sym-
 bolically thus. If $a = b$, and
 $c = b$, then $a = c$, if we deny
 this, we deny that a magnitude

is equivalent to itself. There would be no unity in the diversity of things which is logically necessary to knowledge. A similar argument applies to the other axioms referred to. Such axioms are presupposed in all mathematical reasoning and hence are not purely geometrical.

It is the axioms that are necessary to the science of Geometry that we desire to consider.

If we consider the eighth axiom - magnitudes that can be made to coincide

with the same magnitude are equal," we find that it differs from the first axiom. For when we apply this axiom to geometrical magnitudes it involves the assumption that geometrical magnitudes can be moved from one part of space to another part of space without change of form or dimensions. This assumption is founded on our experience with the physical world.

As, if we consider the eleventh axiom which asserts that through a point not in a straight line only one line

can be drawn parallel to the given line, we shall find that it involves an assumption founded on experience.

Again there are assumptions implied in the definitions and postulates. A definition is the explanation of the meaning of a term or word. Now if this explanation involves terms that are not definable we must assume them. In our definition of the plane we must assume the elementary idea of extension or

space. Proof in Euclidean Geometry consists in general in establishing the congruence of certain lines and angles. And to establish this congruence the lines and the angles are supposed to be moved from one part of space to another. Now the question arises as to whether the nature of space might not interfere with this movement of geometrical figures. We can easily imagine surfaces upon which triangles could not be freely moved without

change of form.

And since it is an assumption that through three points not in a right line a plane surface can be drawn, we see that it would be possible for a triangle to lie on a surface which would not admit of free movement to any part of its surface without change of form.

Not every surface has the unique property that figures lying on it can be freely moved from one part to any other part without change of form or dimensions

This special property was first printed out by Gauss in a discussion on distance in which he says that the radius of curvature at every point of the surface must be the same.

Every surface that fulfils this condition admits of the free movement of the figures lying upon it and without change of form or dimensions. There are three surfaces of this kind. The spherical surface which has a positive constant radius of curvature, the plane surface whose radius of curvature

is zero, and (3) the pseudo-spherical surface which has a constant negative radius of curvature. So we see that if we assume that the figures of Geometry can be freely moved from any position to any other position in space without change of form or dimensions, the surface is one of three different kinds of surfaces, the spherical surface, the plane surface or the pseudo-spherical surface.

The eleventh axiom which is known as the "axiom of parallels" has been a subject for much discussion. If we deny this axiom we are led to the Geometry upon the pseudo-spherical surface first given by Lobatschewsky and Bolyai.

If we consider the twelfth axiom
 "Two lines can not enclose a space"
 we see that by making this ass-
 umption we exclude the spher-
 ical surface upon which all lines
 meet.

Thus we see that the last
 or twelfth axiom distinguishes
 the plane surface and pseudo-
 spherical surface geometry
 from the spherical surface geom-
 etry. And that the axiom of
 parallels distinguishes the plane
 surface geometry of Euclid from
 the pseudo-spherical surface
 geometry of Lobatchewsky and
 Bolyai.

And hence we are led to
 conclude that these latter axioms
 is the eighth eleventh and twelfth are

determined by experience and are not necessary to experience of the subject-matter of Geometry. Hence are not a priori axioms.

Now the other axioms of Euclid which are necessary to the analytical mathematics as well as to Geometry are said to be entirely a priori. In considering the foundations of Geometry the natural tendency is to attempt to construct a Geometry based entirely upon these a priori axioms. Such a Geometry would be free from the assumptions founded upon experience. This cannot be done, for Geometry is a physical science and must depend upon some assumptions determin-

led on account of experience. The narrowest approach to this kind of Geometry however is the projective Geometry first given by Pappus in his "Theory of Distances" and since developed by Klein.

We will now consider the axioms of Geometry from the point of view of projective Geometry.

If we assume, to start with, the plane (which is not necessarily the Euclidean plane) and points in the plane, then how are we to distinguish between the different points of the plane? For without this distinction, we could have no Geometry. The existence of a set of points in a plane

is not sufficient. There must be some relation between the points before we can reason about them. And hence the next axiom is necessary is. the straight line. This axiom simply states that for any two points there exists a straight line completely determined by the two points. The line is a unique relation between any two points.

Now there are an infinite number of points on any straight line and since this straight line is the only relation existing between any two points on this line, any set of two points that we may choose is indistinguishable

from any other set of two points on the line. For the relation connecting each set of two points is simply the line.

Points are qualitatively indistinguishable from each other and the points on the line are distinguished by the line relation from the other points of the plane. It follows that the straight line, which is a figure determined by two points which are qualitatively indistinguishable from each other, is qualitatively indistinguishable from any other line of the plane.

If we consider the point as the intersection of two straight

lines, we see that two lines passing through any point are qualitatively indistinguishable from any other two lines passing through the same point, since the point is determined by any two lines passing through it.

Hence we have a basis for the definition of equality in projective Geometry which is independent of the notion of distance.

If we take a point O in the plane and any straight line not through this point and take any four points on this line as A, B, C, D , we have four new

straight lines determined
by these four points i.e. OA ,
 OB , OC , and OD . - see the fig.

Now without changing the
four lines OA , OB , OC and OD , the
four points A , B , C , D could have
been situated at A' , B' , C' , D' or
at the intersections of any
other straight lines with the
four lines OA , OB , OC , and OD .
The points A , A' , A'' - - - - - bring ab-

ways situated on the line OA and the points $B, B', B'' \dots$ always situated on the line OB and in the same way the points $C, C', C'' \dots$ and $D, D', D'' \dots$ are always situated on the lines OC and OD .

The different sets of four four points as described above $A, B, C, D, A', B', C', D', A'', B'', C'', D''$, &c. we define as being equal.

In a similar way we consider the four lines OA, OB, OC , and OD . The point O being qualitatively indistinguishable from any other point of the plane, we define these two sets of four straight lines

are equal if each set of four lines passes through a single point and if the corresponding lines of different sets pass two by two through four points in a straight line.

We are led to these definitions by considering one, two, and three, points in the plane. If we consider one point, it is qualitatively indistinguishable from any other point and we have no ^{projective} method of determining any other point. If we consider two points they determine the straight line passing through them which may be taken as a projective relation to det-

minus a third point. But the third point can lie any place on this line and hence this relation is not sufficient to determine the third point.

However, if we consider three points on a straight line, we have something more than the line relation and the fourth point can be determined.

Suppose that we have any three points on a straight line as A, B, C , (see the figure below) and if we take any point outside the line as O , the three lines OA, OB , and OC are determined. We take any point on OC as M and draw

equal to the set of points L, M, O, C ,
 and the set L, M, O, C equal to the
 set L, N, P, C . also the set of points
 L, N, P, C is equal to the set
 D, B, A, C . Hence we have the set
 of points $A, B, C, D =$ the set of points
 D, B, A, C . And we see that the
 points A, B, C, D can be projectively trans-
 formed into the points
 D, B, A, C . This is an important
 principal and we will refer to
 it again.

Thus we see that, in
 this kind of geometry where
 the notion of distance is not
 used, if we are given
 three points on a line as
 A, B, C , we can uniquely det-

words line and point.

Hence we define the point as the relation determined by two lines, and the line as the relation determined by two points.

We have also defined the manner by which different sets of points are distinguishable from each other and the equality between the figures in this abstract Geometry without the use of quantitative comparisons or the notion of distance.

Points alone are indistinguishable from each other, and with points only we

could have no quantity. Now
 the straight line relation between
 two points gives us a certain
 identity of quality which according
 to Logic is necessary to
 quantitative comparison. Any
 two points are qualitatively in-
 distinguishable from any other
 two points on the line as all
 points on the line are connected
 by the same relation. And
 since the straight line is the re-
 lation between two points, lines
 are only distinguishable from
 each other by means of the
 points. And if only one point
 is given, it is impossible to
 distinguish any difference.

between the lines through
the point.

And we conclude
that all geometrical figures
which are projectively related
are qualitatively indistinguish-
able from each other.

Now we desire to
know the axioms upon
which this Geometry is founded.
In the first place we have
assumed the point, line,
and plane, but the lines
and planes are not ne-
cessarily Euclidean. Next we
assume that space is hom-
ogeneous. This assumption
is absolutely necessary for

if this were not true points would not be qualitatively indistinguishable from each other, and the four points situated on a transversal could not be defined as equal to four points situated on another transversal in another part of space. A point or position is the same in all parts of space and is indefinable aside from its relation to some other position, hence all positions are relative.

Our next assumption or axiom is that there exists a unique relation between

any two points called a straight line, between any three not in a straight line a unique rotation called a plane and so on for a finite number of points.

For, if this were not true, a finite number of relations of a point to a given number of points would not be sufficient to determine its relation to a new point. And since it is only possible for us to consider a finite number of relations, Geometry could not exist. That is our space must have a finite integral

number of dimensions.

These axioms are necessary to any form of externality and since some form of externality is necessary to experience, it follows that these axioms are necessary to experience and hence are a priori axioms.

we will next consider the axioms of metrical geometry in the light of the principles deduced from our considerations of projective geometry.

The new element which enters into our considerations is distance and the idea of measurement.

In projective geometry our comparisons of geometrical figures was purely qualitative. But the idea of measurement implies a quantitative comparison

of magnitudes. We saw in projective Geometry that if we consider one point and any other point that it was necessary to assume some relation as existing between them before we could reason about them. This relation was unique so far as any two points on the line were concerned. Hence we could not distinguish one point from any other particular point. Now this is what is necessary in metrical Geometry. We must have a relation which will enable

us to distinguish one point on a line from any other point on the line. This relation is the distance between ~~the~~ the two points. This is the first axiom of metric Geometry.

And in projective Geometry we found that it was necessary to assume that space was everywhere the same. And in metric Geometry we must assume that space will admit of the free movement of geometrical figures from one part of space

to any other part of space
And our second axiom is
the axiom of free mobility.

In our first axiom
we assume that there exists
a unique relation between
two points is the distance
between the two points. Now
measurement logically involves
the comparison of magnitudes.
It is no measurement of
a distance to divide it
into 100 equal parts. Those
parts must be referred to
other magnitudes, there
is no absolute measure

of distance and in the last analysis we must refer our magnitudes to arbitrarily chosen units of magnitude.

Now when we desire to compare two spatial magnitudes as two distances we assume that our unit of distance can be made to coincide with the first distance a certain number of times and then with the second distance a certain number of times, and the ratio of these two numbers is the ratio of the two distances. And this can

only be done (we assume) if space admits of the movement of the point of reference from one part of space to any other without change of dimensions.

The third axiom in Geometry is the axiom of dimensions. In the projective Geometry we found that we must logically have a finite integral number of dimensions. In metric Geometry we limit the number to three. This is not logically necessary for a position might be as log-

ically determined by four relations as by three, hence we see that the axiom contains an a priori and an empirical element.

The notion of distance being indefinable, we must discuss the method of determining the measure of distance. This can not be done analytically but depends upon experience. Now we will consider the notion from the analytical stand point; and find the possible determinations of the measure of distance

which will not contradict the three necessary axioms of geometry given above.

As a preliminary to the discussion of the measure of distances we define the following functions $f(z)$, $g(z)$, and e^z .

$$f(z) = 1 + \frac{z^2}{12} + \frac{z^4}{14} + \dots$$

$$g(z) = z - \frac{z^3}{13} + \frac{z^5}{15} + \dots$$

$$e^z = 1 + z + \frac{z^2}{2} + \frac{z^3}{6} + \frac{z^4}{24} + \dots$$

The power series representing the above functions are convergent for all values of z , real and imaginary, and hence represent analytic functions of z .

From these three equations we have (2) $e^{iz} = f(z) + i g(z)$, and $f(0) = 1$, also $f(2) = 1 - 2(1 - \frac{2}{3} \frac{2}{4}) - \frac{2^6}{16}(1 - \frac{2}{7} \frac{2}{8}) - \dots$. This is negative, hence we conclude that there is a value between 0 & 2 which will satisfy the equation $f(z) = 0$ we call this value or number $\frac{\pi}{2}$.

In (1) if we substitute $-iz$ for z and iz for z in e^z we have $f(z) = \frac{1}{2}(e^{iz} + e^{-iz})$ (3) and $g(z) = \frac{1}{2i}(e^{iz} - e^{-iz})$ (4). From 3 & 4 we have $f(z)^2 + g(z)^2 = 1$ (5)

And since the number $\frac{\pi}{2}$ made $f(z) = 0$ if we substitute it for z in equation (5),

we have $q\left(\frac{\pi}{2}\right) = \pm 1$. And from (2) we have

$$e^{i\frac{\pi}{2}} = \pm i, (6) \quad \therefore e^{i\pi} = -1.$$

and $e^{2i\pi} = 1$ hence $(e^{(2\pi+i\pi)i})^{\lambda} =$

$$e^{\pi\lambda} = -1. \quad \text{This shows that}$$

the function $e^{\pi i \lambda}$ has a period 2π and from (3) and (4) the functions $f(z)$ and $g(z)$ have the same period.

The function $u = e^z$ is an analytic function of z but we can not say that z is an analytic function of u . For $z = \log \rho + (w + 2n\pi)i$ where $\rho = \text{mod.}$ and $w = \text{arg. of } u$ and $2\pi = \text{the period}$. Now if we define $z = \log \rho + wi$

in which case $n=0$ and if we call this the principal value of $\log$ where w varies from $-\pi$ to $+\pi$, then to each and every value of u there exists one and only one value for z .

We also define the functions $C(z)$, $S(z)$, and $T(z)$, as follows:

$$(8) \begin{cases} 1) C(z) = \frac{1}{2} (e^{\frac{z}{K}} + e^{-\frac{z}{K}}) \\ 2) S(z) = \frac{1}{2} (e^{\frac{z}{K}} - e^{-\frac{z}{K}}) \\ 3) T(z) = \frac{e^{\frac{z}{K}} - e^{-\frac{z}{K}}}{e^{\frac{z}{K}} + e^{-\frac{z}{K}}} \end{cases}$$

From 1 and 2 of equations 8 we have $S(-z) = -S(z)$, $C(-z) = C(z)$ and

$C(x)^2 - S(x)^2 = 1$, also with 3 of equation

$$(8) \quad T(x) = \frac{S(x)}{C(x)} \quad (9)$$

The determination of the measure of magnitude in Geometry can be reduced to two fundamental problems To determine the distance between two points and to determine the difference in direction of two intersecting straight lines.

In order to apply our analytical relations to the simplest spatial relations and in the simplest manner, we assume

the plane and the line and
the point is the plane,
the point not on the line.

And here, as before, the plane
and the line are not ^{necessarily} the Eucli-
dian plane and line. The
point is the intersection of two
straight lines the line is
a relation between two points
and the plane a relation
between three points.

Let l, l' be the lines
p any point outside
the line and p' the line
which makes the angle $po l$ =
the angle $p'o l'$ and let

n be any other point on the line, and let θ represent the

angle θ of n .

If now we represent the measure of the distance on by z or any constant relation to z , then for each and every position of n on the line there is a unique value of $T(z)$ and conversely for every value of $T(z)$ there is a unique

position of n on the line.
 We define $\tan \theta$ to be $\frac{on}{op}$
 since op is constant and
 n is any point on the line,
 for every position of n there
 exists a unique value of
 $\tan \theta$ and conversely for
 every value of $\tan \theta$ there
 exists a unique value of
 on or a unique position of
 n on the line.

Then $\tan \theta$ and $T(n)$
 are functions so related that
 to any value θ of $\tan \theta$ there
 corresponds a unique value

2 of $T(z)$. If we suppose that the line pm turns about the point p it will take different positions on the line and to each position there is a unique θ and z .

If we express the relation between $T(z)$ and $\tan \theta$ in its most general form $A T(z) \tan \theta + B T(z) + C \tan \theta + D = 0$, we see that when pm takes the position p_0 , $m = 0$, $\tan \theta = 0$, and $T(z) = 0$. This shows that $D = 0$.

Next suppose that m takes the position m_i

symmetrical to o in the line ol' . If we call $on = z_0$ then $on' = -z_0$ and our general form then becomes,

$$A T(z) \tan \theta_0 + B T(z) + C \tan \theta_0 = 0 \text{ for } z_0$$

$$\text{and } A T(z) \tan \theta_0 - B T(z) - C \tan \theta_0 = 0 \text{ for } -z_0$$

By adding we have $A = 0$ and by subtracting we have $-\frac{C}{B} = \frac{T(z_0)}{\tan \theta_0}$

And the relation between $T(z)$ and $\tan \theta$ for different values of on takes the form.

$$10) \quad \frac{Tz}{\tan \theta} - \frac{Tz_0}{\tan \theta_0} = -\frac{C}{B} = \text{constant.}$$

$$11) \text{ or } T(z) = \frac{\tan \theta}{\tan \Delta} \text{ where } \tan \Delta = \frac{\tan \theta_0}{T(z_0)}$$

In (3) of equation 8 if we let z become infinite we have

$T(z) = 1$ or if z becomes minus infinity, we have $T(z) = -1$. And from (10) and (11) we have Δ and $-\Delta$ as the values of θ for $z = \infty$ and $-\infty$. Whence we conclude that from any point external to a line we can draw two lines pm_1 and pm_2 meeting the line at ∞ and they form with po an angle Δ given by equation (11).

We can find the value z for any point by solving the equation $\frac{\epsilon^{\frac{z}{k}} - \epsilon^{-\frac{z}{k}}}{\epsilon^{\frac{z}{k}} + \epsilon^{-\frac{z}{k}}} = \frac{\tan \theta}{\tan \Delta}$ which gives for $\theta < \Delta$ the real value

$$Z = \frac{K}{2} \log \frac{\tan \Delta + \tan \theta}{\tan \Delta - \tan \theta} \text{ and for } \theta > 0$$

$$Z = \frac{K}{2} \log \frac{\tan \Delta + \tan \theta}{\tan \Delta - \tan \theta} + \frac{K}{2} \pi i.$$

Hence with our form
 for the measure of distance
 we arrive at the conclusion
 of Lobatchewsky that the
 lines passing through the
 point p fall into two classes
 1 those situated in the angle
 n, p, n_1 and 2 those separated
 by the parallels p, n_1 and
 p, n_1 . The angle Δ is called
 the angle of parallelism for
 the point p . The form (11)

shows that Δ depends upon ϕ which we will call d hence we will assume $\frac{1}{\tan \Delta} = \phi(d)$ and the form (10) becomes

(12) $T(\Delta) = \phi(d) \tan \theta$. Also (11) shows $\tan \theta = 0$ when $T(\Delta) = 0$, but $T(\Delta) = 0$ when $\kappa = \infty$, when $\tan \theta = 0$ $\theta = 0$ and $\tan \Delta = \infty$ or $\Delta = 90^\circ$

whatever be the value of d . And this particular value of κ gives only one parallel through the point p , which corresponds to the parallelism of Euclid. Thus our expression for the distance between

two points contains a variable parameter λ .

If we apply this principle to a triangle ABC in which the angle A corresponds to θ , B to ϕ , and C to ψ , calling the sides a, b, c , we have from (11)

$T(a) = \phi(b) \tan A$, $T(b) = \phi(a) \tan B$ from the first we have $\frac{T_a}{\tan A} = \phi(b)$ and

$$\frac{1}{\tan^2 A} = \frac{\phi(b)}{T(a)}, \quad 1 + \frac{1}{\tan^2 A} = 1 + \frac{\phi(b)}{T(a)},$$

and $\frac{\tan A}{1 + \tan^2 A} = \frac{T(a)}{\sqrt{T(a)^2 + \phi(b)^2}}$, we designate the first member of this equation by $g'(A)$

$$(13) \quad \therefore g'(A) = \frac{T(a)}{\sqrt{T(a)^2 + \phi(b)^2}} \quad \text{Substituting}$$

for $T(a) \frac{S(a)}{C(a)}$ we have

$$(14) \quad g'(A) = \frac{S(a)}{\sqrt{S(a)^2 + \phi(b)^2 + S(a)^2 \phi(b)^2}} \quad \text{and in}$$

the same way we can find $g'(B)$ for the angle B .

$$(15) \quad g'(B) = \frac{T(b)}{\sqrt{T(b)^2 + \phi(a)^2}} = \frac{S(b)}{\sqrt{S(b)^2 + \phi(a)^2 + S(b)^2 \phi(a)^2}} \quad (16)$$

In short form if $g'(A) = 0$ or $g'(B) = 0$ then $a = 0$ and $b = 0$. If $g'(A) = 1$ and $g'(B) = 1$ then $a = c$ and $b = c$

Hence we conclude that ~~that~~
 $g'(A)$ and $g'(B)$ must be of the
 form of $\frac{f'(a)}{f'(c)}$ and $\frac{f'(b)}{f'(c)}$ and we

$$\text{have } \phi(a) = T(a) = f(a), \phi(b) = T(b) = f(b) \quad (17)$$

from 13 and 15, and from 14 and 16

$$\phi(a) = s(a) = f(a) \quad \phi(b) = s(b) = f(b) \quad (18)$$

Again from (11) we have
 $T(a) + \phi(b) = \phi(b)(\tan^2 A + 1)$ and

$$\frac{1}{\tan^2 A + 1} = \frac{\phi(b)}{T(a) + \phi(b)} = f'(A).$$

$$(19) \quad \therefore f'(A) = \frac{\phi(b)}{\sqrt{T(a) + \phi(b)}} = \frac{\phi(b)\sqrt{1 + s^2(a)}}{\sqrt{s^2(a) + \phi(b) + s^2(a)\phi(b)}}.$$

And in the same way we form

$$f'(B) = \frac{\phi(a)}{\sqrt{T(b) + \phi(a)}} = \frac{\phi(a)\sqrt{1 + s^2(b)}}{\sqrt{s^2(b) + \phi(a) + s^2(b)\phi(a)}}.$$

When $f'(A) = 0$ $b = 0$ if $f'(A) = 1$, $b = c$.
 When $f'(B) = 0$ $a = 0$ if $f'(B) = 1$ $a = c$.

And in the same way we have

$$(21) \quad \phi(b) = T(b) = f(b), \quad \phi(a) = T(a) = f(a)$$

$$(22) \quad \phi(b) = S(b) = f(b), \quad \phi(a) = S(a) = f(a).$$

Combining 17 with 11 and 12 we have,

$$23 \quad \left\{ \begin{array}{l} g'A = \frac{T_a}{T_c}, \text{ and } g'(B) = \frac{T(b)}{T(c)}, \\ f'A = \frac{T(b)}{T(c)}, \text{ and } f'(B) = \frac{T(a)}{T(c)}, \\ T(c)^2 = T(a)^2 + T(b)^2. \end{array} \right.$$

Let α and β be the segments of c made by a perpendicular from o upon c then we have

$$T\alpha = T(b) f'A = \frac{T^2(b)}{T(c)}, \text{ and}$$

$$24 \quad T\beta = T(a) f'(B) = \frac{T^2(a)}{T(c)} \text{ adding we}$$

$$\text{have, } T\alpha + T\beta = \frac{T^2(b) + T^2(a)}{T(c)} = Tc = T(\alpha + \beta)$$

This last equation makes $S(\alpha) S(\beta) = 0$

From (8) this becomes

$$\left(\frac{a}{k} + \frac{1}{L^2} \left(\frac{a}{k} \right)^2 + \dots \right) \left(\frac{b}{k} + \frac{1}{L^2} \left(\frac{b}{k} \right)^2 + \dots \right) = 0.$$

For this to be true k must $= \infty$ however for $k = \infty$ the ratio $\frac{T(a)}{T(c)}$ reduces to $\frac{a}{c}$ and $\frac{T(b)}{T(c)} = \frac{b}{c}$.

From 23 we have

$$(23) \quad \begin{cases} g'(A) = \frac{a}{c}, & g'(B) = \frac{b}{c} \\ f'(A) = \frac{b}{c}, & f'(B) = \frac{a}{c} \end{cases}$$

$c^2 = a^2 + b^2$, These are the formulas of plane trigonometry, ($g'(A) \times f'(A)$ being the $\sin A$ and $\cos A$), deduced from the general form by giving to k the value ∞

If we combine (12) with (9) and (10) we have

$$(26) \quad \left\{ \begin{array}{l} g(A) = \frac{S(a)}{S(c)}, \quad g(B) = \frac{S(b)}{S(c)}, \\ f(A) = \frac{T(b)}{T(c)}, \quad f(B) = \frac{T(a)}{T(c)}, \\ c(c) = c(a), \quad c(b) = \frac{1}{\tan A \tan B}, \\ c(a) = \frac{f(A)}{g(A)}, \quad c(b) = \frac{f(B)}{g(B)}, \\ \tan B = \frac{T(a)}{T(b)}, \quad \tan A = \frac{T(b)}{T(a)}. \end{array} \right.$$

Also from (12) we have

$$g(d) = S(d), \quad \tan \Delta = \frac{1}{S(d)},$$

$$g(A) = \frac{1}{c(d)}, \quad f(A) = T(d),$$

Now if we consider any triangle ABC , we will find that it depends upon

the value of k as to whether our triangle represents a Euclidean plane triangle, a pseudo-spherical triangle, or a spherical triangle.

From (5) we have

$$g(A, s(b)) = g(B, s(a)) = s(s),$$

$$\tan C_a = -\frac{1}{\tan A C(b)}, \quad \tan C_b = -\frac{1}{\tan B C(a)}.$$

$$\tan.(C_a + C_p) = -\tan C = \frac{\tan A C_b + \tan B C_a}{1 - \tan A \tan B C_a C_b},$$

$$\tan (C_a + C_p) = -\tan T(c) = \frac{T(a) f(B) + T(b) f(A)}{1 - T(a) T(b) f(A) f(B)}.$$

from which we have

$$\frac{S(a)}{g(A)} = \frac{S(b)}{g(B)} = \frac{S(c)}{g(C)}.$$

$$\tan B \tan A \tan C = \frac{\tan A}{C_a} + \frac{\tan B}{C_b} + \frac{\tan C}{C_a C_b}.$$

$$-T(a) T(b) T(c) = \frac{T(a)}{f(A)} + \frac{T(b)}{f(B)} + \frac{T(c)}{f(A) f(B)}.$$

$$\frac{S^2(a)}{g^2(A)} = \frac{S^2(b)}{g^2(B)} = \frac{S^2(c)}{g^2(C)} = \frac{1 - f(A) - f(B) - f(C) - 2 f(A) f(B) f(C)}{g^2(A) g^2(B) g^2(C)}.$$

$$= \frac{S^2(a) S^2(b) S^2(c)}{1 - C_a^2 - C_b^2 - C_c^2 + 2 \cdot C_a C_b C_c}.$$

27)

$$C_a = C_b C_c + S(b) S(c) f(A)$$

$$C_b = C_a C_c + S(c) S(a) f(B)$$

$$C_c = C_a C_b + S(a) S(b) f(C).$$

$$f(A) = f(B) f(C) - g(B) g(C) C_a.$$

$$f(B) = f(C) f(A) - g(C) g(A) C_b.$$

$$f(C) = f(A) f(B) - g(A) g(B) C_c.$$

The formulas (27) correspond to non-Euclidean Geometry for different values of K .

We have seen that for $K = \infty$ we have the plane triangle of Euclid, if in the formulas of (27), we give to K any real constant value we have the pseudo-spherical triangle and the formulas (27) become the formulas of pseudo-spherical trigonometry.

However, if we let $\frac{1}{K} = i$, we have a spherical triangle and the formulas (27) become

the formulas of spherical trigonometry for example

$$C(a) = C(b)C(c) + S(b)S(c) \cos A, \text{ becomes}$$

$$f(A) = f(b)f(c) - g(b)g(c)f(A) \text{ by virtue}$$

$$\text{of the relation } C(iZ) = f(Z)$$

$$\text{and } S(iZ) = i g(Z) \text{ See Ex. and (4)}$$

$$\text{or } \cos A = \cos b \cos c - \sin b \sin c \cos A, \text{ and}$$

we get similar results from the other forms.

For this value of k our form (10) $\frac{TZ}{\tan \theta} = \text{constant}$, becomes

$$\frac{g(Z)}{f(Z)} = \tan \theta h \text{ where } h = \text{a constant}$$

and for all values of θ , Z is finite. For $\tan \theta = \infty$ we have

$$\frac{g(Z)}{f(Z)} = \infty \text{ and the value of } Z$$

which satisfies this equation is $\frac{\pi}{2}$. In this case the value of Σ has an upper limit and there are no parallels.

Thus with this relation for the distance between two points we are led to three different Geometries or to a general Geometry of which the plane, Spherical, and pseudo-spherical Geometries, are particular cases.

In the immediate neighborhood of a point these forms become indistinguishable from the Euclidean Geometry.

For if we replace $S(x)$ and $C(x)$ by their values,

$$S(x) = \frac{x}{K} + \frac{1}{12} \frac{x^3}{K^3} + \dots$$

$$C(x) = 1 - \frac{1}{2} \frac{x^2}{K^2} + \dots$$

And make x small enough that x^2 and higher powers of x may be neglected which is equivalent to reducing the sides of our triangle to very short distances, we have

$S(x) = \frac{x}{K}$ and $C(x) = 1$. Now if we substitute these values in (27), we have

$$\frac{\frac{a}{K}}{\sin A} = \frac{\frac{b}{K}}{\sin B} = \frac{\frac{c}{K}}{\sin C} = \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}.$$

$$\tan A \tan B \tan C = \tan A + \tan B + \tan C.$$

$$1 - \cos^2 A - \cos^2 B - \cos^2 C - 2 \cos A \cos B \cos C = 0.$$

$$a^2 = b^2 + c^2 + 2bc \cos A.$$

$$\cos A = \cos (B + C).$$

In these latter forms we have placed $\cos z$ and $\sin z$ for $f(z)$ and $g(z)$, and these are formulas of plane trigonometry.

Thus we conclude that there are three axioms necessary to the existence of Geometry.

And that with these three axioms we can logically construct a general Geometry in which our expression for the distance between

two points contains an arbitrary constant K . And if we allow K to take all possible values we obtain three different forms of Geometry as we have seen.

And in order to distinguish between these different forms of Geometry, it is necessary to make other assumptions which depend entirely upon experience. For we have seen that if our experience is confined to the neighborhood of a point it is impossible to

distinguish between the three forms.

We learn from experience that a line in our actual space has no upper limit and that two lines never enclose a space and this excludes the spherical form of Geometry. We also learn from experience that only one ^{line} parallel ^{to a given} line can be drawn through a point outside the given line, and this distinguishes the pseudo-spherical form from the Euclidean form of Geometry. And

hence these axioms,
are wholly empirical
axioms.

I approve this thesis
for the Master's degree.

Mr. Glodney's work this
year in Mathematics has
been good.

J. N. Fellows.

RECEIVED
OCT 9 1913
UNIV. OF MO.

118654

University of Missouri - Columbia

010-100725789

213

~~This book is not to be taken from the library
Neither is it to be loaned out overnight.~~

Reserve Book R.R. - 205
Permanent Collection