

# EXTENSION OF WEAKLY AND STRONGLY $F$ -REGULAR RINGS BY FLAT MAPS

IAN M. ABERBACH

## §1. INTRODUCTION

Throughout this paper all rings will be Noetherian of positive characteristic  $p$ . Hence tight closure theory [HH1–4] takes a prominent place (see §2 for tight closure definitions and terminology). The purpose of this note is to help answer the following question: if  $R$  is weakly (resp. strongly)  $F$ -regular and  $\phi : R \rightarrow S$  is a flat map then under what conditions on the fibers is  $S$  weakly (resp. strongly)  $F$ -regular. This question (among many others) is raised in [HH4] in section 7. It is shown there that if  $\phi$  is a flat map of local rings,  $S$  is excellent and the generic and closed fibers are regular then weak  $F$ -regularity of  $R$  implies that of  $S$  (Theorem 7.24). One of our main results weakens the hypotheses considerably.

**Theorem 3.4.** *Let  $\phi : (R, \mathfrak{m}) \rightarrow (S, \mathfrak{n})$  be a flat map. Assume that  $S/\mathfrak{m}S$  is Gorenstein and  $R$  is weakly  $F$ -regular and Cohen-Macaulay. Suppose that either*

- (1)  $c \in R^\circ$  is a common test element for  $R$  and  $S$ , and  $S/\mathfrak{m}S$  is  $F$ -injective, or
- (2)  $c \in S - \mathfrak{m}S$  is a test element for  $S$  and  $S/\mathfrak{m}S$  is  $F$ -rational, or
- (3)  $R$  is excellent and  $S/\mathfrak{m}S$  is  $F$ -rational.

*Then  $S$  is weakly  $F$ -regular.*

We note that the Gorenstein assumption on the fiber is essential, even if  $R$  is regular. Even weakening the assumption on the fiber to  $\mathbb{Q}$ -Gorenstein is not strong enough to give a good theorem, as Singh [Si] gives an example of  $R \rightarrow S$  flat, where  $R$  is a discrete valuation domain,  $S/\mathfrak{m}S$  is  $\mathbb{Q}$ -Gorenstein and strongly  $F$ -regular, yet  $S$  is not weakly  $F$ -regular!

We also prove a corresponding result for strong  $F$ -regularity.

**Theorem 3.6.** *Let  $(R, \mathfrak{m}, K) \rightarrow (S, \mathfrak{n}, L)$  be a flat map of  $F$ -finite reduced rings with Gorenstein closed fiber. Assume that  $R$  is strongly  $F$ -regular. If  $S/\mathfrak{m}S$  is  $F$ -rational then  $S$  is strongly  $F$ -regular.*

In order to prove the first of these theorems we investigate how flat maps  $\phi : (R, \mathfrak{m}) \rightarrow (S, \mathfrak{n})$  with Gorenstein closed fibers affect tight closure for  $I \subseteq R$  such that  $l(R/I) < \infty$

---

1991 *Mathematics Subject Classification.* 13A35.

The author was partially supported by the NSF.

and  $I$  is irreducible in  $R$ . In general these results do *not* depend on the relationship of  $R/\mathfrak{m} \rightarrow S/\mathfrak{n}$  (e.g., separability or finiteness).

While not directly relevant to this paper, we note that other authors have recently investigated tight closure properties under good flat maps. For instance Enescu [En] and Hashimoto [Ha] have recently shown that for a flat map with  $F$ -rational base and  $F$ -rational closed fiber, the target is  $F$ -rational (in the presence of a common test element).

## §2. BACKGROUND FOR TIGHT CLOSURE

Let  $R$  be a Noetherian ring of characteristic  $p > 0$ . We use  $q = p^e$  for a varying power of  $p$  and for an ideal  $I \subseteq R$  we let  $I^{[q]} = (i^q : i \in I)$ . Also let  $R^\circ$  be the complement in  $R$  of the union of the minimal primes of  $R$ . Then  $x$  is in the *tight closure* of  $I$  if and only if there exists  $c \in R^\circ$  such that  $cx^q \in I^{[q]}$  for all  $q \gg 0$ . If  $I^* = I$  then  $I$  is said to be *tightly closed*. We will say that  $I$  is *Frobenius closed* if  $x^q \in I^{[q]}$  for some  $q$  always implies that  $x \in I$ .

There is a tight closure operation for a submodule  $N \subseteq M$ , but we will not discuss this case in general. It is however useful to discuss tight closure in the case of a particular type of direct limit. Suppose that  $M = \varinjlim_t R/I_t$  for a sequence of ideals  $\{I_t\}$ . Let  $u \in M$  be an element which is given by  $\{u_t\}$  where in the direct limit system  $u_t \mapsto u_{t+1}$ . We will say that  $u \in 0_M^*$  if there exists  $c \in R^\circ$  and a sequence  $t_q$  such that for all  $q \gg 0$ ,  $cu_{t_q}^q \in I_{t_q}^{[q]}$ . We will say that  $u$  is in the *finitistic tight closure* of 0 in  $M$ ,  $0_M^{*fg}$ , if there exists  $c \in R^\circ$  and  $t > 0$  such that  $cu_t^q \in I_t^{[q]}$  for all  $q$ . This definition of finitistic tight closure agrees with that in [HH2] for this case. Clearly  $0_M^{*fg} \subseteq 0_M^*$ .

A ring  $R$  in which every ideal is tightly closed is called *weakly  $F$ -regular*. If every localization of  $R$  is weakly  $F$ -regular then  $R$  is  *$F$ -regular*. When  $R$  is reduced then  $R^{1/p}$  denotes the ring of  $p$ th roots of elements of  $R$ . More generally,  $R^{1/q}$  is the ring of  $q$ th roots. Clearly  $R \subseteq R^{1/q}$ . If  $R$  is  $F$ -finite and reduced ( $R^{1/p}$  is a finite  $R$ -module) then  $R$  is called *strongly  $F$ -regular* if for all  $c \in R^\circ$ , there exists a  $q$  such that the inclusion  $Rc^{1/q} \subseteq R^{1/q}$  splits over  $R$ . If  $R$  is  $F$ -finite and  $R_c$  is strongly  $F$ -regular for some  $c \in R^\circ$ , then  $R$  is strongly  $F$ -regular if and only if there exists  $q$  such that  $Rc^{1/q} \subseteq R^{1/q}$  splits over  $R$  [HH1, Theorem 3.3]. Strongly  $F$ -regular rings are  $F$ -regular, and weakly  $F$ -regular rings are normal and under mild conditions (e.g., excellent) are Cohen-Macaulay.

The equivalence of the three conditions is an important open question. Let  $(R, \mathfrak{m})$  be an excellent reduced local ring and let  $E$  be an injective hull of the residue field of  $R$ . Then  $E$  can be written as a direct limit of the form above since  $R$  is approximately Gorenstein. Weak  $F$ -regularity of  $R$  is equivalent to  $0_E^{*fg} = 0$  [HH2, Theorem 8.23], while strong  $F$ -regularity is equivalent to ( $F$ -finiteness and)  $0_E^* = 0$  [LS, Proposition 2.9].

By a *parameter ideal* in  $(R, \mathfrak{m})$  we mean an ideal generated by part of a system of parameters. We say that  $(R, \mathfrak{m})$  is  *$F$ -rational* if every parameter ideal is tightly closed, and  *$F$ -injective* if every parameter ideal is Frobenius closed (this is a slightly different notion of  $F$ -injectivity from that in [FW], but is equivalent for CM rings).  $F$ -rational rings are normal and under mild conditions are Cohen-Macaulay. In a Gorenstein ring,  $F$ -rationality is equivalent to all forms of  $F$ -regularity.

A priori, the multiplier element  $c$  in the definition of tight closure depends on both  $I$  and  $x$ . If  $c$  works for every tight closure test then we say that  $c$  is a *test element* for  $R$ .

If  $c$  works for every tight closure test for every completion of every localization of  $R$  then we say that  $c$  is a *completely stable test element*. It is shown in [HH4] that if  $(R, \mathbf{m})$  is a reduced excellent domain,  $c \in R^\circ$ , and  $R_c$  is Gorenstein and weakly  $F$ -regular then  $c$  has a power which is a completely stable test element for  $R$ .

In [HH2, HH3] it is shown that the multiplier  $c$  in the definition of tight closure need not remain constant. Let  $R$  be a domain. One may have a sequence of elements  $c_q$  such that  $c_q x^q \in I^{[q]}$  where  $c_q$  must have “small order.” We can obtain a notion of order, denoted  $\text{ord}$ , by taking a  $\mathbb{Z}$ -valued valuation on  $R$  which is non-negative on  $R$  and positive on  $\mathbf{m}$ . Let  $R^+$  be the integral closure of  $R$  in an algebraic closure of the fraction field of  $R$  ( $R^+$  has many wonderful properties, such as being a big Cohen-Macaulay algebra for  $R$  when  $R$  is excellent [HH5]). The valuation then extends to a function on  $R^+$  which takes values in  $\mathbb{Q}$ . In particular,  $\text{ord}(c^{1/q}) = \text{ord}(c)/q$ . We will need to use the following theorem [HH3, Theorem 3.1].

**Theorem 2.1.** *Let  $(R, \mathbf{m})$  be a complete local domain of characteristic  $p$ , let  $x \in R$  and let  $I \subseteq R$ . Then  $x \in I^*$  if and only if there exists a sequence of elements  $\epsilon_n \in (R^+)^\circ$  such that  $\text{ord}(\epsilon_n) \rightarrow 0$  as  $n \rightarrow \infty$  and  $\epsilon_n x \in IR^+$ .*

In fact we would like to strengthen this theorem in order to apply it to tight closure calculations for non finitely generated modules which are defined by a direct limit system of ideals. The proof we give is just an altered version of the proof of Theorem 3.1 given in [HH3]. The key component is [HH3, Theorem 3.3]:

**Theorem 2.2.** *Let  $(R, \mathbf{m}, k)$  be a complete local domain. Let  $\text{ord}$  be a  $\mathbb{Q}$ -valued valuation on  $R^+$  nonnegative on  $R$  (and hence on  $R^+$ ) and positive on  $\mathbf{m}$  (and, hence, on  $\mathbf{m}^+$ ). Then there exists a fixed real number  $\nu > 0$  and a fixed positive integer  $r$  such that for every element  $u$  of  $R^+$  of order  $< \nu$  there is an  $R$ -linear map  $\phi : R^+ \rightarrow R$  such that  $\phi(u) \notin \mathbf{m}^r$ .*

The generalization of Theorem 2.1 is given below.

**Theorem 2.3.** *Let  $(R, \mathbf{m})$  be a complete local domain of characteristic  $p$ . Let  $M = \varinjlim_t R/I_t$  be an  $R$ -module and let  $x \in M$ . Suppose that  $x$  comes from the sequence  $\{x_t\}$  where  $x_t \mapsto x_{t+1}$ . Then  $x \in 0_M^*$  if and only if there exists a sequence of elements  $\epsilon_n \in (R^+)^\circ$  such that  $\text{ord}(\epsilon_n) \rightarrow 0$  as  $n \rightarrow \infty$  and for each  $n$  there exists  $t$  such that  $\epsilon_n x_t \in I_t R^+$ .*

*Proof.* The “only if” part is trivial, as if  $cx^q = 0$  for all  $q \gg 0$  then we can take  $\epsilon_q = c^{1/q}$ .

To see the “if” direction, choose  $\nu > 0$  and  $r$  as in Theorem 2.2. Fix  $q = p^e > 0$ . Choose  $n$  large enough that  $\text{ord}(\epsilon_n) < \nu/q$ . Let  $\epsilon = \epsilon_n^q$ . Then there exists  $t$  such that  $\epsilon x_t^q \in I_t^{[q]} R^+$  and  $\text{ord}(\epsilon) < \nu$ . Applying an  $R$  linear map  $\phi$  as in Theorem 2.2 we find that  $c_q x_t^q \in I_t^{[q]} \subseteq (I_t^{[q]})^*$  with  $c_q = \phi(\epsilon) \in R - \mathbf{m}^r$ . Thus, setting  $J_q = \cup_t (I_t^{[q]})^* :_R x_t^q$  we have  $c_q \in J_q$  for all  $q$ .

The sequence  $J_q$  is nonincreasing. If for some  $t$ ,  $yx_t^{pq} \in (I_t^{[pq]})^*$  then  $c'(yx_t^{pq})^{q'} \in (I_t^{[pq]}[q'])^{[q']} = (I_t^{[pqq']})^{[q']}$  for all  $q' \gg 0$  where  $c' \neq 0$ . But then  $c'(yx_t^q)^{pq'} \in (I_t^{[q]}[pq'])^{[q']}$  for all  $q' \gg 0$  and hence  $yx_t^q \in (I_t^{[q]})^*$ , as required.

Since the sequence  $\{J_q\}_q$  is nonincreasing, it cannot have intersection 0, or Chevalley's theorem would give  $J_q \subseteq \mathfrak{m}^r$  for  $q \gg 0$ . As  $c_q \in J_q - \mathfrak{m}^r$  for all  $q$ , we can choose a nonzero element  $d \in \bigcap_q J_q$ . Then for each  $q$  there exists  $t$  such that  $dx_t^q \in (I_t^{[q]})^*$ . If  $c$  is a test element for  $R$  then  $cdx_t^q \in I_t^{[q]}$ . Thus  $x \in 0_M^*$ .  $\square$

**Proposition 2.4.** *Let  $(R, \mathfrak{m})$  be an excellent local domain such that its completion is a domain. Let  $M = \varinjlim_t R/I_t$  be a direct limit system. Fix  $u \notin 0_M^*$ . Then there exists  $q_0$  such that  $J_q = \bigcup_q (I_t^{[q]} : u_t^q) \subseteq \mathfrak{m}^{[q/q_0]}$  for all  $q \gg 0$  (where  $\{u_t\}$  represents  $u \in M$  and  $u_t \mapsto u_{t+1}$ ). In particular if  $I \subseteq R$  we may take  $M = R/I$  where the limit system consists of equalities. Then  $u \notin I^*$  implies that  $(I^{[q]} : u^q) \subseteq \mathfrak{m}^{[q/q_0]}$ .*

*Proof.* Suppose that we can show that the proposition holds in  $\widehat{R}$ . Then  $(I_t^{[q]} :_R u_t^q) \subseteq (I_t^{[q]} :_{\widehat{R}} u_t^q) \cap R \subseteq \mathfrak{m}^{[q/q_0]} \widehat{R} \cap R \subseteq \mathfrak{m}^{[q/q_0]} R$ . Thus we may assume that  $R$  is complete.

For  $x \in R$  let  $f(x)$  be the largest power of  $\mathfrak{m}$  that  $x$  is in, and set  $\mathbf{f}(x) = \lim_{n \rightarrow \infty} f(x^n)/n$ . By the valuation theorem [Re, Theorem 4.16], there exist a finite number of  $\mathbb{Z}$ -valued valuations  $v_1, \dots, v_k$  on  $R$  which are non-negative on  $R$  and positive on  $\mathfrak{m}$  and positive rational numbers  $e_1, \dots, e_k$  such that  $\mathbf{f}(x) = \min\{v_i(x)/e_i\}$ . Furthermore, since  $R$  is analytically unramified, there exists a constant  $L$  such that for all  $x \in R$ ,  $f(x) \leq \lfloor \mathbf{f}(x) \rfloor \leq f(x) + L$  ([Re, Theorem 5.32 and 4.16]).

Now, by Theorem 2.3, for each  $v_i$  there exists a positive real number  $\alpha_i$  such that if  $c \in (I_t^{[q]} : u_t^q)$  then  $v_i(c) \geq \alpha_i q$ . Combined with the valuation theorem we see that  $\mathbf{f}(c) \geq \min\{q\alpha_i/e_i\}$ . Let  $\alpha = \min\{\alpha_i/e_i\}$ . Then  $f(c) \geq \alpha q - L - 1$ . Let  $s = \mu(\mathfrak{m})$ . Choose  $q_1 > 1/\alpha$ ,  $q_2 \geq L + 1$ , and  $q_3 \geq s$  (all powers of  $p$ ). Set  $q_0 = q_1 q_2 q_3$ . Then  $f(c) \geq \alpha q - (L + 1) \geq q/q_1 - (L + 1) \geq q/q_1 q_2 - 1 \geq (q/q_0)s - 1$ . A simple combinatorial argument shows that  $\mathfrak{m}^{(q/q_0)s-1} \in \mathfrak{m}^{[q/q_0]}$ . Hence  $c \in \mathfrak{m}^{[q/q_0]}$ .  $\square$

### §3. TIGHT CLOSURE IN FLAT EXTENSION MAPS

We show in this section that extending a weakly (respectively, strongly)  $F$ -regular ring by a flat map with sufficiently nice Gorenstein closed fiber yields another weakly (resp., strongly)  $F$ -regular ring. These results are Theorems 3.4 and 3.6 (see also Corollary 3.5 for the  $F$ -regular case).

By saying that  $\phi : (R, \mathfrak{m}) \rightarrow (S, \mathfrak{n})$  is flat we mean that  $\phi$  is flat and that  $\phi(\mathfrak{m}) \subseteq \mathfrak{n}$ . Since the map is flat we then know that given ideals  $A, B \subseteq R$  we have  $AS :_S BS = (A :_R B)S$  ( $B$  finitely generated). The next lemma merely asserts that modding out by elements which are regular in the closed fiber preserves flatness.

**Lemma 3.1.** *Let  $\phi : (R, \mathfrak{m}) \rightarrow (S, \mathfrak{n})$  be a flat map. Let  $z_1, \dots, z_d \in S$  be elements whose images in  $S/\mathfrak{m}S$  are a regular sequence. Then for any ideal  $I$  generated by monomials in the  $z$ 's, the ring  $S/IS$  is flat over  $R$ .*

*Proof.* See, for example [HH4, Theorem 7.10a,b].  $\square$

The next proposition shows that tight closure behaves well for irreducible  $\mathfrak{m}$ -primary ideals when extending to  $S$ . Given a sequence of elements  $\mathbf{z} = z_1, \dots, z_d$  we will use  $\mathbf{z}^{[t]}$  to denote  $z_1^t, \dots, z_d^t$ .

**Proposition 3.2.** *Let  $\phi : (R, \mathfrak{m}, K) \rightarrow (S, \mathfrak{n}, L)$  be a flat map with Gorenstein closed fiber. Let  $\mathbf{z} = z_1, \dots, z_d \in S$  be elements whose images form a s.o.p. in  $S/\mathfrak{m}S$ . Let  $I \subseteq R$  be such that  $l(R/I) < \infty$  and  $\dim_K(0 :_{R/I} \mathfrak{m}) = 1$ . Suppose that either*

- (1)  *$R$  and  $S$  have a common test element and  $S/\mathfrak{m}S$  is  $F$ -injective, or*
- (2)  *$c \in S - \mathfrak{m}S$  is a test element for  $S$ , and  $S/\mathfrak{m}S$  is  $F$ -rational, or*
- (3)  *$R$  is excellent,  $\widehat{R}$  is a domain, and  $S/\mathfrak{m}S$  is  $F$ -rational.*

*Then  $I$  is tightly closed in  $R \iff$  for all  $t > 0$ ,  $IS + (\mathbf{z})^{[t]}S$  is tightly closed in  $S \iff$  there exists  $t > 0$  such that  $IS + (\mathbf{z})^{[t]}S$  is tightly closed in  $S$ .*

*Proof.* Let  $b \in S$  have as its image the socle element in  $S/\mathfrak{m}S + (\mathbf{z})S$ . Let  $u \in R$  be the socle element mod  $I$ . Then the socle element of  $S/(IS + (\mathbf{z})S)$  is  $ub$  since the map  $R/I \rightarrow R/I \otimes S = S/IS$  is flat with Gorenstein fibers (there is only one fiber).

Suppose that  $I$  is tightly closed. There is no loss of generality in taking  $t = 1$ . If  $IS + (\mathbf{z})S$  is not tightly closed in  $S$  then we have  $c(ub)^q \in (I^{[q]} + (\mathbf{z})^{[q]})S$  for all  $q$ . In case (1) we may take  $c \in R^\circ$ , so that

$$b^q \in (I^{[q]} + (\mathbf{z})^{[q]})S :_S cu^q = (I^{[q]} :_R cu^q)S + (\mathbf{z})^{[q]}S \subseteq \mathfrak{m}S + (\mathbf{z})^{[q]}S$$

for all  $q \gg 0$ . The first equality is a consequence of flatness, while the inclusion follows since  $u \notin I^*$ . By our assumption that  $S/\mathfrak{m}S$  is  $F$ -injective we reach the contradictory conclusion that  $b \in ((\mathbf{z}) + \mathfrak{m})S$ . In case (2) we have

$$cb^q \in (I^{[q]} + (\mathbf{z})^{[q]})S :_S u^q = (I^{[q]} :_R u^q)S + (\mathbf{z})^{[q]}S \subseteq \mathfrak{m}S + (\mathbf{z})^{[q]}S$$

for all  $q \gg 0$ . As  $S/\mathfrak{m}S$  is  $F$ -rational, it is a domain, so  $c \neq 0$  in  $S/\mathfrak{m}S$ . This contradicts our hypothesis that  $S/\mathfrak{m}S$  is  $F$ -rational (in fact it is enough to assume that  $I$  is Frobenius closed to reach this conclusion). In case (3) we can choose  $q_0$  as in Proposition 2.4, and then

$$c(b^{q_0})^{q/q_0} \in (I^{[q]} + (\mathbf{z})^{[q]})S :_S u^q = (I^{[q]} :_R u^q)S + (\mathbf{z})^{[q]}S \subseteq \mathfrak{m}^{[q/q_0]}S + ((\mathbf{z})^{[q_0]})^{[q/q_0]}$$

for all  $q/q_0$ . But then  $b^{q_0} \in (\mathfrak{m}S + (\mathbf{z})^{[q_0]})^*$ . By persistence, the image of  $b^{q_0}$  is in  $((\mathbf{z})^{[q_0]}S/\mathfrak{m}S)^*$ , which contradicts the  $F$ -rationality of  $S/\mathfrak{m}S$ .

Suppose now that  $IS + (\mathbf{z})^{[t]}S$  is tightly closed in  $S$  for all  $t$ , but  $I$  is not tightly closed in  $R$ . Then  $u \in (IR)^* \subseteq (I + (\mathbf{z})^{[t]})^*$  (since  $R^\circ \subseteq S^0$ ). But then  $u \in \bigcap_t (IS + (\mathbf{z})^{[t]}S)^* \cap R \subseteq \bigcap_t (IS + (\mathbf{z})^{[t]}S) \cap R \subseteq IS \cap R = IR$ .

Finally, suppose that  $(IS + (\mathbf{z})^{[t_0]}S)$  is tightly closed for some  $t_0$ . Given any  $t$ , the socle element of  $(IS + (\mathbf{z})^{[t]}S)$  is  $(z_1 \cdots z_d)^{t-1}ub$ . If  $c((z_1 \cdots z_d)^{t-1}ub)^q \in (IS + (\mathbf{z})^{[t]})^{[q]}$  then by flatness,  $c((z_1 \cdots z_d)^{t_0-1}ub)^q \in (IS + (\mathbf{z})^{[t_0]})^{[q]}$ . Therefore, one such ideal tightly closed shows that all such ideals are tightly closed.  $\square$

To deal with strong  $F$ -regularity we need to give a similar proposition with  $R/I$  replaced by the injective hull  $E_R(R/\mathfrak{m})$ . Suppose that we can write  $E = E_R(R/\mathfrak{m}) = \varinjlim_t R/J_t$ , the set  $\{u_t\} \subseteq R$  is a collection of elements such that  $u_t \mapsto u_{t+1}$  in the map  $R/J_t \rightarrow R/J_{t+1}$  and the image of each  $u_t$  in  $E$  is the socle element of  $E$ . It suffices that  $R$  be approximately Gorenstein [Ho2] (e.g., excellent and normal, or even reduced) to obtain  $E$  in this manner. In particular an  $F$ -finite ring is excellent [Ku], so a reduced  $F$ -finite ring is approximately Gorenstein.

**Proposition 3.3.** *Let  $(R, \mathbf{m}, K) \rightarrow (S, \mathbf{n}, L)$  be a flat map of  $F$ -finite reduced rings with Gorenstein closed fiber.*

- (1) *If  $Rc^{1/q} \subseteq R^{1/q}$  splits for some  $q$  (over  $R$ ) and  $S/\mathbf{m}S$  is  $F$ -injective then  $Sc^{1/q} \subseteq S^{1/q}$  splits for some  $q$  (over  $S$ ).*
- (2) *If  $0$  is Frobenius closed in  $E_R(K)$ ,  $S/\mathbf{m}S$  is  $F$ -rational and  $c \in S - \mathbf{m}S$  then there exists  $q$  such that  $Sc^{1/q} \subseteq S^{1/q}$  splits (over  $S$ ).*

*Proof.* Choose  $\mathbf{z} = z_1, \dots, z_d \in S$  elements which generate a s.o.p. in  $S/\mathbf{m}S$ . By [HH4, Lemma 7.10] we have  $E_S(L) = \varinjlim_v S/(\mathbf{z}^{[v]}) \otimes_R E_R(K) = \varinjlim_{v,t} S/(\mathbf{z}^{[v]}) \otimes_R R/J_t = \varinjlim_t S/(\mathbf{z}^{[t]}, J_t)S$ . If  $b \in S$  generates the socle element in  $S/(\mathbf{m} + (\mathbf{z}))S$  then the image of  $(z_1 \cdots z_d)^{t-1}bu_t$  in  $S/((\mathbf{z}^{[t]}) + J_t)S$  maps to the socle element of  $E_S$  (where  $u_t$  is as given above).

In case (1), if for all  $q$  the inclusion  $Sc^{1/q} \rightarrow S^{1/q}$  fails to split, by [Ho1, Theorem 1 and Remark 2] for all  $q$  there exists  $t_q$  such that

$$c(z_1 \cdots z_d)^{(t_q-1)qb^q u_{t_q}^q} \in ((\mathbf{z}^{[t_q]}), J_{t_q})^{[q]}S.$$

Hence  $(z_1 \cdots z_d)^{(t_q-1)qb^q} \in ((\mathbf{z}), J_{t_q})^{[q]} :_S cu_{t_q}^q \subseteq (J_{t_q}^{[q]} :_R cu_{t_q}^q)S + (\mathbf{z}^{[t_q]})^{[q]}S \subseteq \mathbf{m}S + (\mathbf{z}^{[t_q]})^{[q]}S$  for  $q \gg 0$  (we are using here that if  $Rc^{1/q} \subseteq R^{1/q}$  splits for some  $q$  then  $Rc^{1/q'} \subseteq R^{1/q'}$  splits for all  $q' \geq q$ ). Thus  $b^q \in \mathbf{m}S + (\mathbf{z})^{[q]}$  since  $S/\mathbf{m}S$  is CM. This contradicts the  $F$ -injectivity of  $S/\mathbf{m}S$ .

To see (2), if there is no splitting we obtain

$$c(z_1 \cdots z_d)^{(t_q-1)qb^q} \in (\mathbf{z}^{[t_q]}, J_{t_q})^{[q]} :_S u_{t_q}^q \subseteq (J_{t_q}^{[q]} :_R u_{t_q}^q)S + ((\mathbf{z}^{[t_q]})^{[q]}S \subseteq \mathbf{m}S + ((\mathbf{z}^{[t_q]})^{[q]}S$$

and hence  $cb^q \in \mathbf{m}S + (\mathbf{z})^{[q]}$ . This contradicts the  $F$ -rationality of  $S/\mathbf{m}S$ .  $\square$

We can now give our main theorems on the extension of weakly and strongly  $F$ -regular rings by flat maps with Gorenstein closed fiber.

**Theorem 3.4.** *Let  $\phi : (R, \mathbf{m}) \rightarrow (S, \mathbf{n})$  be a flat map. Assume that  $S/\mathbf{m}S$  is Gorenstein and  $R$  is weakly  $F$ -regular and CM. Suppose that either*

- (1)  *$c \in R^\circ$  is a common test element for  $R$  and  $S$ , and  $S/\mathbf{m}S$  is  $F$ -injective, or*
- (2)  *$c \in S - \mathbf{m}S$  is a test element for  $S$  and  $S/\mathbf{m}S$  is  $F$ -rational, or*
- (3)  *$R$  is excellent and  $S/\mathbf{m}S$  is  $F$ -rational.*

*Then  $S$  is weakly  $F$ -regular.*

*Proof.* To see that  $S$  is weakly  $F$ -regular it suffices to show that there exists a sequence of irreducible tightly closed ideals of  $S$  cofinite with the powers of  $\mathbf{n}$ . As  $R$  is weakly  $F$ -regular (so normal) and CM it is approximately Gorenstein. Say that  $\{J_t\}$  is a sequence of irreducible ideals cofinite with the powers of  $\mathbf{m}$ . Let  $\mathbf{z} = z_1, \dots, z_d \in S$  be elements which form a s.o.p. in  $S/\mathbf{m}S$ . Then  $(J_t + \mathbf{z}^{[t]})S$  is a sequence of irreducible ideals in  $S$  cofinal with the powers of  $\mathbf{n}$ . By Proposition 3.2, in cases (1), (2), and (3), the ideals  $(J_t + \mathbf{z}^{[t]})S$  are tightly closed in  $S$  (in case (3),  $\widehat{R}$  is still weakly  $F$ -regular, so is a domain).

Therefore  $S$  is weakly  $F$ -regular. We note that in case (2) we may weaken the assumption that  $R$  is weakly  $F$ -regular to the assumption that  $R$  is  $F$ -pure (see the comment in the proof of Proposition 3.2, part (2)).  $\square$

The next corollary should be compared with [HH4, Theorem 7.25(c)].

**Corollary 3.5.** *Let  $(R, \mathfrak{m}) \rightarrow (S, \mathfrak{n})$  be a flat map of excellent rings with Gorenstein fibers. Suppose that the generic fiber is  $F$ -rational and all other fibers are  $F$ -injective. If  $R$  is  $F$ -regular then  $S$  is  $F$ -regular.*

*Proof.* By hypothesis the generic fiber is Gorenstein and  $F$ -rational, therefore there is a  $c \in R^\circ$  which is a common completely stable test element.  $F$ -regularity is local on the prime ideals of  $S$  and the fiber of such a localization is the localization of a fiber, hence Gorenstein and  $F$ -injective (the property of  $F$ -injectivity is easily seen to localize). Therefore Theorem 3.4(1) always applies.  $\square$

**Theorem 3.6.** *Let  $(R, \mathfrak{m}, K) \rightarrow (S, \mathfrak{n}, L)$  be a flat map of  $F$ -finite reduced rings with Gorenstein closed fiber. Assume that  $R$  is strongly  $F$ -regular. If  $S/\mathfrak{m}S$  is  $F$ -rational then  $S$  is strongly  $F$ -regular.*

*Proof.* We must show that there exists an element  $c \in S^0$  such that  $S_c$  is strongly  $F$ -regular and  $S_c^{1/q} \subseteq S^{1/q}$  splits for some  $q$ .

If there exists  $c \in R^\circ$  such that  $S_c$  is strongly  $F$ -regular (i.e., a power of  $c$  is a common test element for  $R$  and  $S$ ) then we are done by Proposition 3.3(1). Even if  $R$  and  $S$  have no (apparent) common test element, however, we claim that there exists  $c \in S - \mathfrak{m}S$  such that  $S_c$  is strongly  $F$ -regular. Once we have shown this, the theorem follows by Proposition 3.3(2).

Since the non-strongly  $F$ -regular locus is closed [HH1, Theorem 3.3] it suffices to show that  $S_{\mathfrak{m}S}$  is strongly  $F$ -regular, for then there exists an element  $c \in S - \mathfrak{m}S$  such that  $S_c$  is strongly  $F$ -regular. Let  $B = S_{\mathfrak{m}S}$ . Then  $R \rightarrow B$  is flat and the closed fiber is a field. In particular  $E_B(B/\mathfrak{m}B) = E_R(K) \otimes_R B$ . As  $R$  is strongly  $F$ -regular (so normal) it is approximately Gorenstein. Say  $E_R = \varinjlim_t R/J_t$  with socle element mapped to by  $u_t$  (as before). Then  $u_t \in B/J_t B$  will still map to the socle element  $u$  in  $E_B$ . Suppose that  $u \in 0_{E_B}^*$ . This means there exists  $b \in B_0$  such that for all  $q$  there exists  $t_q$  such that  $bu_{t_q}^q \in J_{t_q}^{[q]} B$ . Hence  $b \in J_{t_q}^{[q]} :_B u_{t_q}^q = (J_{t_q}^{[q]} :_R u_{t_q}^q) B$ . Note that  $R$  is an excellent normal domain, so its completion remains a domain. Thus by Proposition 2.4 we see that as  $q \rightarrow \infty$ ,  $(J_{t_q}^{[q]} :_R u_{t_q}^q)$  gets into larger and larger powers of the maximal ideal, since 0 is tightly closed in  $E_R$ . Thus  $b \in \bigcap_N \mathfrak{m}^N B = 0$ , a contradiction.  $\square$

## REFERENCES

- [En] F. Enescu, *On the behavior of  $F$ -rational rings under flat base change*, J. of Alg. (to appear).
- [FW] R. Fedder and K.I. Watanabe, *A characterization of  $F$ -regularity in terms of  $F$ -purity*, Commutative Algebra, MSRI Publications No. 15, Springer-Verlag, 1989, pp. 227–245.
- [Ha] M. Hashimoto, *Relative Frobenius maps and Cohen-Macaulay  $F$ -injective homomorphisms*, preprint.
- [Ho1] M. Hochster, *Contracted ideals from integral extensions of regular rings*, Nagoya Math. J. **51** (1973), 25–43.
- [Ho2] M. Hochster, *Cyclic purity versus purity in excellent Noetherian rings*, Trans. A.M.S. **231** (1977), 463–488.

- [HH1] M. Hochster and C. Huneke, *Tight closure and strong  $F$ -regularity*, Memoires Soc. Math. de France **38** (1989), 119–133.
- [HH2] M. Hochster and C. Huneke, *Tight closure, invariant theory, and the Briançon-Skoda theorem*, J. Amer. Math. Soc. **3** (1990), 31–116.
- [HH3] M. Hochster and C. Huneke, *Tight closure and elements of small order in integral extensions*, J. Pure Appl. Alg. **71** (1991), 233–247.
- [HH4] M. Hochster and C. Huneke,  *$F$ -regularity, test elements, and smooth base change*, Trans. Amer. Math. Soc. **346** (1994), 1–62.
- [HH5] M. Hochster and C. Huneke, *Infinite integral extensions and big Cohen-Macaulay algebras*, Annals of Math. **135** (1992), 53–89.
- [Ku] E. Kunz, *On Noetherian rings of characteristic  $p$* , Amer. J. Math **98** (1976), 999–1013.
- [LS] G. Lyubeznik and K. E. Smith, *On the commutation of the test ideal with localization and completion*, Trans. A.M.S (to appear).
- [Re] D. Rees, *Lectures on the asymptotic theory of ideals*, LMS Lecture Note Series 113, Cambridge University Press, Cambridge.
- [Si] A. K. Singh,  *$F$ -regularity does not deform*, Amer. Jour. Math. **121** (1999), 919–929.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA, MO 65211  
*E-mail address:* `aberbach@math.missouri.edu`