

Public Abstract

First Name:William

Middle Name:Gerald

Last Name:Alexander

Adviser's First Name:Patrick

Adviser's Last Name:Shiu

Co-Adviser's First Name:

Co-Adviser's Last Name:

Graduation Term:SP 2011

Department:Biological Sciences

Degree:PhD

Title:Meiotic Silencing by Unpaired DNA: Elucidation of a Post-Transcriptional Gene Silencing Pathway

Meiotic silencing by unpaired DNA is an RNAi-mediated pathway that functions to silence unpaired genes during the sexual phase of *Neurospora crassa*. The presence of a gene insertion or deletion causes an unpairing event during the homolog pairing stage. My dissertation's aim was to further characterize this new RNAi pathway. First, I was involved in adapting the in vivo protein interaction technique Bimolecular Fluorescence Complementation (BiFC) to *N. crassa* to investigate SAD-1/SAD-2 interaction. SAD-1 and SAD-2 interact in the perinuclear region, suggesting that SAD-2 functions to bring SAD-1 to its proper location. Next, I was involved in identifying components of quelling that also function in meiotic silencing. *dcl-1* and *qip* were found to be important for sexual development and meiotic silencing. DCL-1 and QIP also localize in the perinuclear region, implicating these genes in meiotic silencing. Finally, I tested the mating type locus for immunity to meiotic silencing. Unpaired reporter gene insertions resulted in ascospore populations exhibiting partial protection from silencing. When a reporter gene was enclosed with paired mat locus flanks at an ectopic locus, partial protection was again observed.