

Jessica Struckhoff

Major: Biology

University: University of Missouri-Columbia

Faculty Mentor: Dr. Mark D. Kirk

Mentor Department: Biological Sciences

Funded by: Life Sciences Undergraduate Research Opportunity Program

The effect of Camgaroo-2 incorporation on the differentiation potential of embryonic stem cells

Jessica Struckhoff, Katie Spears, Chris Pierret and Mark D. Kirk

Embryonic stem (ES) cells are capable of differentiating into any cell type in the body and are a promising therapeutic agent. Our research focuses on the differentiation of ES cells into functional neurons and/or glial that can nurture host cells of the nervous system that are damaged due to disease. Cells must express the appropriate phenotype and perform the proper function after transplantation. Camgaroo-2 is a fluorescence protein that provides a basal fluorescence and responds to a rise in intracellular calcium by producing an increase in fluorescence emission. Our lab transfected a mouse ES cell line (GSI-1) with the Camgaroo-2 gene and is testing this fluorescence indicator to determine the physiological function of cells grown in vitro. There is concern that the incorporation of the Camgaroo-2 gene could alter the cell phenotype, potentially decreasing their differentiation potential. GSI-1 cells were plated on culture slides following a neural induction protocol that uses retinoic acid and allowed to proliferate. Immunohistochemistry of slides was performed to label for neural precursors, immature and mature neurons, astrocytes, and oligodendrocytes (anti-O4). GSI-1 labeling was compared to corresponding immunohistochemistry performed on another ES cell line that had also been 'neuralized' to determine if the differentiation potential of the GSI-1 cells was similar to that of the other ES cell line. Similar labeling was seen for all markers except O4 which did not label for the GSI-1 cells, indicating the GSI-1 cells have the potential to differentiate into all cells of neural lineage except possibly oligodendrocytes. GSI-1 cells retained the ability to differentiate post-transfection with the Camgaroo-2 gene. Because of their unique ability to respond to an influx of intracellular calcium, GSI-1 cells expressing Camgaroo-2 can be transplanted into rodent models for human disease, and can be tested post transplantation for their ability to function as neural cells.