



Roselyn Jan W.
Clemente-Fuentes, MD;
Heather Pickett, DO
Nellis Family Medicine
Residency, Nellis Air Force
Base, Nev

Misty Carney, MLIS
Stimson Library, Fort Sam
Houston, Tex

ASSISTANT EDITOR
Paul Crawford, MD
Nellis Family Medicine
Residency, Nellis Air Force
Base, Nev

Q / How can pregnant women safely relieve low-back pain?

EVIDENCE-BASED ANSWER

A / ACETAMINOPHEN IS SAFE for use in pregnancy but lacks evidence of efficacy (strength of recommendation [SOR]: C, usual practice).

Both physical therapy and water aerobics reduce sick days caused by low-back pain (strength of recommendation [SOR]: B, randomized controlled trial [RCT]). Acupuncture, including auricular acupuncture, also relieves low-back pain and improves function (SOR: B, 2 RCTs).

Osteopathic manipulative therapy

(OMT) slightly improves disability (SOR: B, RCT).

Corticosteroid injection at the sacrospinous ligament insertion decreases pain (SOR: B, RCT).

Insufficient evidence of efficacy exists for support garments (SOR: B, systematic review).

No serious maternal or fetal adverse effects have been reported with any of these therapies.

Evidence summary

Even though clinical research is lacking, acetaminophen is widely used to relieve low-back pain with no documented teratogenic effect (US Food and Drug Administration [FDA] category B). Nonsteroidal anti-inflammatory drugs are classified as FDA category D in the third trimester because of their documented association with oligohydramnios, premature closure of the ductus arteriosus, nephrotoxicity, and periventricular hemorrhage in the fetus.¹ Opioids are category C and a poor choice to treat low-back pain in pregnancy.²

Physical therapy and water aerobics relieve pain, reduce sick days

A 2007 Cochrane review of interventions for treating back pain in pregnancy analyzed 8 studies with a total of 1305 patients that examined the effects of adding physical therapy and acupuncture to usual care.³ In one RCT, 407 patients with and without pain received 5 30-minute individualized physical therapy exercise sessions, 2 45-minute group physical therapy classes, or standard care.^{3,4} Low-back pain decreased with group physical therapy

($P < .05$; number needed to treat [NNT]=3.2) and individual therapy (NNT=2.1). Patients who received individual therapy had a 12% decrease in sick days.

A prospective trial of 258 patients, half of whom did water aerobics and half physical therapy, showed comparable results for the 2 interventions (NNT=11.4 for decreased sick days; odds ratio=0.38, 95% confidence interval [CI], 0.16-0.88).³

Acupuncture reduces pain and analgesic use

A prospective, randomized open study cited in the 2007 Cochrane review divided 72 patients at 24 to 37 weeks' gestational age into a group that received acupuncture plus standard care and a standard-care-only control group.^{3,5} Treatment sessions occurred one or 2 times per week until delivery or recovery. The acupuncture group reported decreased pain (60% vs 14% for controls; $P < .01$; NNT=2.2) and improved function (43% vs 9% for controls; $P < .001$; NNT=2.9). There was also a difference in analgesic use: 0% for the acupuncture group vs 14% for controls; $P < .05$; NNT=7.1.

CONTINUED ON PAGE 268

CONTINUED FROM PAGE 260

A 2009 RCT divided 159 patients at 25 to 38 weeks' gestational age into 3 groups: auricular acupuncture at specific points for one week, sham auricular acupuncture at nonspecific points for one week, and controls. At the end of Week 1, 80% of the acupuncture group had a clinically significant reduction in pain compared with 56% in the sham acupuncture group and 36% in the control group ($P=.001$ acupuncture vs sham, $NNT=4.2$; $P<.0001$ acupuncture vs controls, $NNT=2.3$).⁶

Osteopathic manipulative therapy (OMT) decreases disability, but not pain

A 2010 RCT divided 144 third trimester patients into 3 groups that received usual obstetric care, sham ultrasound therapy plus usual obstetric care, or OMT.⁷ Pain remained similar among the 3 groups throughout the study. Using the 24-point Roland-Morris Disability Questionnaire, OMT decreased disability by 0.72 points (95% CI, 0.31-1.14; $P<.001$) compared with 0.35 points in the usual obstetric-care-only group (95% CI, -0.06 to 0.76; $P=.09$). Ultrasound had no effect.

Corticosteroid injection reduces pain in a small trial

A small RCT of injection with the corticosteroid triamcinolone at the sacrospinous ligament insertion in 36 women with low-back pain showed significant reduction in pain in

17 of 18 women in the triamcinolone group compared with 9 of 18 women in the control group ($P<.01$; $NNT=2$).⁸

Evidence lacking on maternity support garments

A poor-quality systematic review of 10 studies ($N=1909$) of maternity support garments found insufficient evidence because of the heterogeneity of the trials.⁹

Recommendations

The American College of Obstetricians and Gynecologists suggests the following measures to prevent and treat low-back pain in pregnancy:¹⁰

- wear low-heeled (not flat) shoes with good arch support
- get help when lifting heavy objects
- place one foot on a stool or box when standing for long periods
- place a board between the mattress and box spring if the bed is too soft
- squat down, bend knees, and keep back straight when lifting
- sit in chairs with good back support or use a small pillow to provide support
- sleep on side with pillows between knees for support
- apply heat, cold, or massage to the painful area. JFP

 **Acetaminophen is safe for use in pregnancy but lacks evidence of efficacy.**

References

1. Black RA, Hill DA. Over-the-counter medications in pregnancy. *Am Fam Physician*. 2003;67:2517-2524.
2. Vermani E, Mittal R, Weeks A. Pelvic girdle pain and low back pain in pregnancy: a review. *Pain Pract*. 2010;10:60-71.
3. Pennick VE, Young G. Interventions for preventing and treating pelvic and back pain in pregnancy. *Cochrane Database Syst Rev*. 2007;(2):CD001139.
4. Ostgaard HC, Zetherstrom G, Roos-Hansson E, et al. Reduction of back and posterior pelvic pain in pregnancy. *Spine*. 1994;19:894-900.
5. Kvorning N, Holmberg C, Grennert L, et al. Acupuncture relieves pelvic and low back pain in late pregnancy. *Acta Obstet Gynecol Scand*. 2004;83:246-250.
6. Wang SM, Dezinno P, Lin EC, et al. Auricular acupuncture as a treatment for pregnant women who have low back and posterior pelvic pain: a pilot study. *Am J Obstet Gynecol*. 2009;201:271.e1-271.e9.
7. Licciardone JC, Buchanan S, Hensel KL, et al. Osteopathic manipulative treatment of back pain and related symptoms during pregnancy: a randomized controlled trial. *Am J Obstet Gynecol*. 2010;202:43.e1-43.e8.
8. Torstensson T, Lindgren A, Kristiansson P. Corticosteroid injection treatment to the ischiadic spine reduced pain in women with long-lasting sacral low back pain with onset during pregnancy: a randomized, double blind, controlled trial. *Spine*. 2009;34:2254-2258.
9. Ho SS, Yu WW, Lao TT, et al. Effectiveness of maternity support belts in reducing low back pain during pregnancy: a review. *J Clin Nurs*. 2009;18:1523-1532.
10. American College of Obstetricians and Gynecologists. Patient education guidelines for easing back pain during pregnancy. August 2011. Available at: <http://www.acog.org/-/media/For%20Patients/faq115.pdf?dmc=1&ts=20130118T1434071958>. Accessed December 10, 2012.



Visit us @ jfponline.com

THE JOURNAL OF
**FAMILY
PRACTICE**