

Public Abstract

First Name:Hesam

Middle Name:

Last Name:Oveys

Adviser's First Name:Stephen

Adviser's Last Name:Montgomery-Smith

Co-Adviser's First Name:

Co-Adviser's Last Name:

Graduation Term:SP 2015

Department:Mathematics

Degree:PhD

Title:Age-dependent Branching Processes and Applications to the Luria-Delbrck Experiment

Microbial populations adapt to their environment by acquiring advantageous mutations, but in the early twentieth century, questions about how these organisms acquire mutations arose. The experiment of Salvador Luria and Max Delbrück that won them a Nobel Prize in 1969 confirmed that mutations don't occur out of necessity, but instead can occur many generations before there is a selective advantage, and thus organisms follow Darwinian evolution instead of Lamarckian. Since then, new areas of research involving microbial evolution has spawned as a result of their experiment. Determining the mutation rate of a cell is one such area. Probability distributions that determine the number of mutants in a large population have been derived by D. E. Lea, C. A. Coulson, and J. B. S. Haldane. However, not much work has been done when time of cell division is dependent on the cell age, and even less so when cell division is asymmetric, which is the case in most microbial populations. Using probability generating function methods, we rigorously construct a probability distribution for the cell population size given a life-span distribution for both mother and daughter cells, and then determine its asymptotic growth rate. We use this to construct a probability distribution for the number of mutants in a large cell population, which can be used with likelihood methods to estimate the cell mutation rate.