

Public Abstract

First Name:Mary

Middle Name:Elizabeth

Last Name:Heffernan

Adviser's First Name:Catherine

Adviser's Last Name:Peterson

Co-Adviser's First Name:

Co-Adviser's Last Name:

Graduation Term:FS 2007

Department:Nutrition Area Program

Degree:MS

Title:The Effects of Regular Tanning Bed Use and Increased Vitamin D Status on Bone Mineral Density and Serum Inflammatory Markers in Healthy Women

Vitamin D is synthesized in the skin in response to UVB radiation and has an essential function in optimal bone health. Recent evidence has also implicated a role for vitamin D in a properly functioning immune system. The objective of this study was to determine the relationship between vitamin D status, inflammatory markers, and bone mineral density in healthy pre- and post-menopausal women who regularly use tanning beds. This observational study examined 69 healthy female subjects: 49 Tanners and 20 Non-Tanners. Subjects provided medical and dietary information, a blood specimen, and bone mineral density was measured. Blood specimens were analyzed for serum 25(OH)D, parathyroid hormone (iPTH), estradiol (E2), cortisol, and inflammatory markers. Results: Tanners had serum 25(OH)D concentrations that were significantly higher ($P < 0.0001$) and iPTH concentrations that were significantly lower ($P < 0.0001$) than Non-Tanners. There were no differences in bone density between groups. Tanners had significantly lower serum TNF α ($P < 0.0200$) and a linear regression revealed that 25(OH)D had a significant inverse relationship with TNF α ($P < 0.0463$), which remained significant after controlling for potential covariates. Conclusions: Serum 25(OH)D status is inversely related to TNF α concentrations in healthy women, which may in part explain its role in the prevention and treatment of numerous diseases.