

# REGULATION OF PHOSPHORYLATION OF DOPAMINE D3 RECEPTORS IN MOUSE STRIATAL NEURONS *in vivo*

D. Rivera, MD<sup>1</sup>; L.M. Mao, PhD<sup>2</sup>; E.E. Fibuch, MD<sup>1</sup>; J.Q. Wang, PhD, MD<sup>1,2</sup>

<sup>1</sup>Department of Anesthesiology, University of Missouri-Kansas City School of Medicine, Saint Luke's Hospital;  
<sup>2</sup>Department of Basic Medical Science, University of Missouri-Kansas City School of Medicine, Kansas City, Missouri

## INTRODUCTION

- Dopamine D3 receptors (D3Rs) are G-protein-coupled receptors.
- These D3Rs inhibit adenylyl cyclase and the downstream formation of cAMP. Due to their preferential expression in the mesolimbic areas, especially in the nucleus accumbens (NAc), they are known to play a major role in the mesolimbic function.
- Recently, we found that D3Rs are phosphorylated at serine 229 by Ca<sup>2+</sup>/calmodulin-dependent protein kinase II (CaMKII).
- This phosphorylation is subject to the modulation, and the phosphorylation level controls receptor function.
- In this study, an effort was made to develop a phospho- and site-specific antibody.
- Using this antibody, we **hypothesized** that the modulation of D3R phosphorylation at Ser229 by dopamine D1 receptor signals would occur in the mouse NAc *in vivo*.


## METHODS

- To detect phosphorylation of D3Rs at Ser229 in striatal neurons *in vivo*, we developed a phospho- and site-specific antibody.
- A short peptide containing phospho-Ser229 (KRILTRQNpSQICISI) was synthesized and used as an immunogen for producing a polyclonal antibody from rabbits.
- Upon demonstration of the selectivity of this antibody, we used it in Western blot to monitor changes in Ser229 phosphorylation in striatal neurons in response to dopamine D1 receptor stimulation by a D1 selective agonist SKF81297.
- Following IACUC approval, adult male mice (CL57/BL6) were randomly divided into 2 groups (n = 3 per group). These animals received an intraperitoneal injection of saline or SKF81297 (1 mg/kg).
- Animals were sacrificed by cervical dislocation 15 min after drug injection. The ventral striatum (NAc) was removed and homogenized.
- Homogenates were centrifuged (1000 g, 10 min), and the supernatant was used for Western blot.
- Densities of immunoblots were measured using optical scanning and the data were analyzed using Student's t-test (p < 0.05)


## RESULTS

- A series of control experiments validated the selectivity of the antibody against phosphorylated D3Rs at Ser229 (pD3R-Ser229).
- Using this antibody, we found that the level of pD3R-Ser229 in the NAc was significantly increased in mice treated with a systemic injection of the D1 receptor agonist SKF81297 (1 mg/kg, 15 min) compared to mice treated with saline.

Figure 1: Selectivity of the Antibody




L1: Standard  
L2: Phospho-immunogen peptide:  
CKRILTRQNpSQICISI  
L3: Non-phospho-peptide:  
CKRILTRQNSQCISI  
L4: L2 + pre-absorbed antibody



L1: NAc from WT mice  
L2: NAc from WT mice  
L3: NAc from D3R -/- mice  
L4: NAc from D3R -/- mice

Figure 2: Stimulation of dopamine D1 receptors increases phosphorylation of D3Rs in the NAc



## CONCLUSIONS

- A phospho- and site-specific antibody against phospho-D3Rs at Ser229 has seemingly been produced.
- The use of this antibody has revealed that stimulation of dopamine D1 receptors increases phosphorylation of D3Rs in the mouse NAc *in vivo*.

## DISCUSSION

- Our efforts in this study seem to lead to a production of such an antibody. Indeed, using this antibody, we found that Ser229 phosphorylation of D3Rs is subject to the modulation by dopamine D1 receptors.
- This indicates a previously-unrecognized D1-dependent negative feedback control of D3R function given that Ser229 phosphorylation inhibits D3Rs.
- In a subpopulation of striatal neurons that co-express D3Rs and D1 receptors, dopamine is able to concurrently enhance the D1-mediated phosphorylation of D3Rs to induce a heterologous desensitization of D3Rs after their activation.
- This regulation is deemed important for maintaining normal homeostasis of D3R function and could be altered to contribute to the pathogenesis of various neurological disorders.