

6. WHO guidelines on hand hygiene in health care: first global patient safety challenge. Clean care is safer care. World Health Organization - 2009
7. O'Grady NP, Alexander M, Burns LA, Dellinger EP, Garland J, Heard SO, Lipsett PA, Masur H, Mermel LA, Pearson ML, Raad II, Randolph AG, Rupp ME, Saint S; Healthcare Infection Control Practices Advisory Committee (HICPAC). Guidelines for the prevention of intravascular catheter-related infections. *Clin Infect Dis*. 2011 May;52(9):e162-93.
8. Tablan OC, Anderson LJ, Besser R, Bridges C, Hajjeh R; CDC; Healthcare Infection Control Practices Advisory Committee. Guidelines for preventing health-care--associated pneumonia, 2003: recommendations of CDC and the Healthcare Infection Control Practices Advisory Committee. *MMWR Recomm Rep*. 2004 Mar 26;53(RR-3):1-36.
9. Resar R, Pronovost P, Haraden C, Simmonds T, Rainey T, Nolan T. Using a bundle approach to improve ventilator care processes and reduce ventilator-associated pneumonia. *Jt Comm J Qual Patient Saf*. 2005 May;31(5):243-8.
10. Hooton TM, Bradley SF, Cardenas DD, Colgan R, Geerlings SE, Rice JC, Saint S, Schaeffer AJ, Tambayh PA, Tenke P, Nicolle LE; Infectious Diseases Society of America. Diagnosis, prevention, and treatment of catheter-associated urinary tract infection in adults: 2009 International Clinical Practice Guidelines from the Infectious Diseases Society of America. *Clin Infect Dis*. 2010 Mar 1;50(5):625-63.

A CASE OF DRESS SYNDROME DUE TO VANCOMYCIN

Danielle Zimmerman, M4, William Steinmann, MD

University of Missouri – Columbia

BACKGROUND

Drug reaction with eosinophilia and systemic symptoms (DRESS) syndrome is a severe systemic reaction that usually begins 2-6 weeks after the introduction of the inciting agent. It is most commonly characterized by fever, rash, hematologic abnormalities (eosinophilia, atypical lymphocytosis), lymphadenopathy, HHV-6 reactivation and internal organ involvement. At least 10% of the cases are fatal. Treatment includes withdrawal of the offending medication and administration of corticosteroids, though the efficacy of the latter has not been fully evaluated.

CASE REPORT

A 64 year-old white male with past medical history of type 2 diabetes mellitus, hypertension, hyperlipidemia, COPD, coronary artery disease, gout, and a fifteen pack-year smoking history presented with complaints of fever, rash and abdominal discomfort. Patient developed a new pruritic, nontender rash on back, chest, arms associated with bilateral arm swelling and shortness of breath. Eight months prior to the admission, he had an open reduction for fracture of the left distal femur after a motor vehicle accident. He had two admissions related to this. One month previously, he was admitted with left septic knee, which was treated by arthrocentesis, screw removal, irrigation and drainage. Intravenous vancomycin was given and vancomycin beads were placed in the knee joint space. Eight days prior to the arrival, the patient was readmitted with complaints of fever, chills, left knee pain and the X-ray of the left knee was suggestive of septic arthritis, which was subsequently managed with irrigation and drainage. Vancomycin was continued and the patient was discharged two days prior to the current admission. The patient hadn't had any recent medication changes and denied any exposure to exotic pets, molds, dusts, chemicals, and drugs.

On physical exam, his temperature was 37.8 °C with a blood pressure of 98/64mmHg. The exam

was remarkable for bilateral upper extremity edema and limited range of motion in the left knee secondary to mild swelling, without erythema or tenderness to palpation or movement. The skin exam revealed a very subtle, erythematous, papular, non-tender and non-confluent pruritic rash on chest and abdomen, more prominent on back and arms. Nikolsky sign was negative.




Figure A: Back showing rash with hyperemia and blanching; hospital day 6

Figure B: Edematous fingers and rash, left knee s/p ORIF in background; hospital day 6

Laboratory results on admission revealed white blood cell count of $10,300/\mu\text{L}$, with 16.5% eosinophil's (normal 0-10%), and platelet count of $390,000/\mu\text{L}$. Other labs included alkaline phosphatase of 332 U/L, AST of 225 U/L, and ALT of 413 U/L compared to previous normal LFT's. Urinalysis was unremarkable and opacities, likely due to atelectasis. Peripheral smear from later hospital course showed large atypical eosinophil's and no toxic granulations in neutrophils. Blood culture remained negative during the hospital stay. Chest x-ray and abdominal ultrasound were unremarkable.

Given the cutaneous changes, eosinophilia and other systemic signs, a presumptive diagnosis of DRESS syndrome was made. Vancomycin was discontinued and patient was switched to Daptomycin. Prednisone 60 mg daily was initiated for treatment of DRESS syndrome. The white counts increased to $18,800/\mu\text{L}$ on day 2, so steroids were held with concern regarding its immunosuppressive effects. Allopurinol was discontinued on hospital day 3, given its reported association with DRESS syndrome. However, as shown in the table over the next several days the white blood cell count escalated dramatically with accompanying eosinophilia. Without further signs of infection in the left knee, prednisone 60 mg was restarted on hospital day 7 to treat presumptive DRESS syndrome.

The prednisone was tapered from 60mg to 30mg daily on hospital day 9. (The patient's white cell count peaked at $44,000/\mu\text{L}$ on hospital day 10, and then decreased substantially by the day of discharge). Improvement of the patient's rash and edema was noted starting on hospital day 3 and continued until discharge, when the rash was no longer visible. However, bilateral axillary lymphadenopathy persisted. (HHV-6 titer drawn on hospital day 2 were negative; however, a repeat HHV-6 titer drawn on hospital day 8 came back positive). Due to absence of left shift and

Progression of WBC Count				
Date	Hospital Day	WBC X 10 ³ / μL	% Eosinophils	Treatment Notes
8/13	2	10.3	16	Daptomycin begun One dose prednisone given
8/14	3	18.8	5	Daptomycin increased Allopurinol discontinued
8/15	4	19.9	26	
8/16	5	22.7	20	
8/17	6	29.1	21	
8/18	7	29.9	18	Prednisone restarted 60 mg
8/19	8	29.9	22	Prednisone 60 mg
8/20	9	36.5	22	Prednisone 30 mg
8/21	10	44.0	10	Prednisone 30 mg
8/22	11	38.7	32	Prednisone 30 mg
8/23	12	33.3	41	Prednisone 30 mg
8/24	13	22.1	39	Prednisone 30 mg
8/25	14	15.6	31	Prednisone 30 mg

toxic granulations in the neutrophils the infection and steroid response were less likely. Clonal hypereosinophilic disorder was also deemed unlikely because the eosinophils had been normal within the month. Because the patient continued to have complaints of left knee pain and given the leukocytosis, daptomycin was continued during this hospitalization. Patient was discharged on Prednisone 30mg for total of ten days, and then tapered down to 10 mg every 10 days.

DISCUSSION

DRESS syndrome was originally identified as a hypersensitivity reaction to certain anticonvulsants, including carbamazepine, lamotrigine, and phenobarbital, but a number of other medications with aromatic groups have been reported to cause the syndrome (1, 2, 3, and 5), including about 18 cases reported with allopurinol and four with vancomycin (2). Our patient had been on allopurinol for several years and he had started vancomycin about four weeks prior to the onset of symptoms, which makes vancomycin the most likely causative agent. More support for this conclusion includes the score of 4 which is a “possible adverse drug reaction” according to Naranjo Adverse drug Reaction Scale (1) and a “definite” case of DRESS syndrome per the criteria described by Kardaun et. al (6). Our patient also met all criteria for DRESS syndrome using the diagnostic criteria established by a Japanese consensus group (7).

This case illustrates the challenge in making a definitive diagnosis of DRESS syndrome. As most of the clinical features of DRESS syndrome are non-specific, it may be difficult to exclude other conditions. In this case, our patient’s history of recent surgery and infection, leukocytosis, and leg pain made it difficult to rule out re-infection of the knee. Though infection of the knee remained a concern, DRESS syndrome was considered as highly probable because of the patient’s multiple negative blood cultures and absence of other signs of infection other than knee pain. The most common manifestations of DRESS syndrome include fever, rash, and systemic symptoms, including lymphadenopathy, leukocytosis, and organ damage. Our patient had elevated liver enzymes, axillary adenopathy and bilateral arm edema.

RegiSCAR Diagnostic Criteria (6)	Naranjo Adverse Drug Reaction Scale
Hospitalization Reaction suspected to be drug-related Acute rash Fever > 38 °C* Enlarged lymph nodes at a minimum of two sites* Involvement of at least one internal organ* Blood count abnormalities* Lymphocytes above or below normal limits Eosinophils above the laboratory limits Platelets below the laboratory limits *3/4 of these are required to make the diagnosis	1. Are there previous conclusive reports on this reaction? <i>Yes (+1) No (0) Do not know or not done (0)</i> 2. Did the adverse event appear after the suspected drug was given? <i>Yes (+2) No (-1) Do not know or not done (0)</i> 3. Did the adverse reaction improve when the drug was discontinued or a specific antagonist was given? <i>Yes (+1) No (0) Do not know or not done (0)</i> 4. Did the adverse reaction appear when the drug was readministered? <i>Yes (+2) No (-1) Do not know or not done (0)</i> 5. Are there alternative causes that could have caused the reaction? <i>Yes (-1) No (+2) Do not know or not done (0)</i> 6. Did the reaction reappear when a placebo was given? <i>Yes (-1) No (+1) Do not know or not done (0)</i> 7. Was the drug detected in any body fluid in toxic concentrations? <i>Yes (+1) No (0) Do not know or not done (0)</i> 8. Was the reaction more severe when the dose was increased, or less severe when the dose was decreased? <i>Yes (+1) No (0) Do not know or not done (0)</i> 9. Did the patient have a similar reaction to the same or similar drugs in any previous exposure? <i>Yes (+1) No (0) Do not know or not done (0)</i> 10. Was the adverse event confirmed by any objective evidence? <i>Yes (+1) No (0) Do not know or not done (0)</i> Scoring > 9 = definite ADR 5-8 = probable ADR 1-4 = possible ADR 0 = doubtful ADR

The pathogenesis of DRESS syndrome is complex and has yet to be completely elucidated. Both drug metabolism and genetic susceptibility have been implicated. Hapten-carrier adduct formation, by either the drugs themselves or their metabolites, and noncovalent drug presentation (8) may lead to T cell activation with an overrepresentation of CD8⁺ cells. It has been hypothesized that the increased risk for DRESS syndrome and other drug reactions among patients with certain HLA variants is due to the affinity of certain drugs, their metabolites, or hapten-carrier adducts for specific HLA binding sites.

These activated T cells produce the cytokines TNF α , IFN γ , and IL-2 that mediate the “cytokine storm,” which may be responsible for the wide constellation of symptoms seen in DRESS syndrome (8, 9). The expansion and activation of CD8⁺ cells may cause the reactivation of latent human herpes viruses seen at the onset of DRESS syndrome, typically HHV-6, but also HHV-7, CMV, EBV, and VZV (9). The broad inflammatory response in DRESS syndrome encourages expansion of regulatory T cells, which are susceptible to infection by viruses such as HHV-6. The infection and compromise of regulatory T cells may mediate the abnormal immunological function observed in this condition. While it has been proposed that polymorphisms in genes for detoxification enzymes may also contribute to DRESS syndrome by increasing the concentration of immunologically active or toxic metabolites, no such polymorphisms have yet been found (8).

Detection of HHV-6 DNA and anti-HHV-6 IgG by PCR analysis is considered sensitive and specific for the diagnosis of DRESS syndrome outside of the United States (10). Thymus and activation-regulated Chemokine (TARC/CCL17), a chemokine, has recently been proposed as an early marker for DRESS syndrome in a study of 29 patients, due to its role in T_H2 type immunity (11).

Current therapy is limited to withdrawal of the offending agent and initiation of corticosteroids to temper the immune response. Our patient received oral prednisone, and the extreme leukocytosis began to subside four days after continuous therapy. Furthermore, his rash and edema had dissipated by the time of discharge. However, it should be noted that there have been no controlled trials validating steroid therapy for this condition (1, 2, and 4). There are several case reports of DRESS syndrome which showed improvement with IVIG, but given the expense and lack of evidence, it is used on an experimental basis only (12, 13). While there are not yet any definitive indicators of mortality, a recent publication showed that an erythema multiforme-like rash, as opposed to the other cutaneous phenotypes possible with DRESS syndrome, may be prognostic for worse hepatic function during the course of illness (14). Sequelae may include end-organ dysfunction (13) and autoimmune disease (15, 16). The latter may be due to viral reactivation and deregulated immune response (16).

Physicians should maintain a low threshold of suspicion for DRESS syndrome when their patients present with a fever, rash, leukocytosis and eosinophilia and a history of new medication use. Further study of documented cases of DRESS syndrome would be helpful in gaining a better understanding of the pathophysiology and optimal therapy for this potentially life threatening disorder.

REFERENCES

1. Ganeva M, Gancheva T, Lazarova R, Troeva J, Baldaranov I, Vassilev I, Hristakieva E, Tzaneva V. "Carbamazepine-induced drug reaction with eosinophilia and systemic symptoms (DRESS) syndrome: report of four cases and brief review." *Int J Dermatol*. 2008 Aug; 47(8):853-60.
2. Cacoub P, Musette P, Descamps V, Meyer O, Speirs C, Finzi L, Roujeau JC. "The DRESS syndrome: a literature review." *Am J Med*. 2011 Jul; 124(7):588-97. Epub 2011 May 17.
3. Fleming P, Marik PE. "The DRESS syndrome: the great clinical mimicker." *Pharmacotherapy*. 2011 Mar; 31(3):332.
4. Walsh SA, Creamer D. "Drug reaction with eosinophilia and systemic symptoms (DRESS): a clinical update and review of current thinking." *Clin Exp Dermatol*. 2011 Jan; 36(1):6-11. doi: 10.1111/j.1365-2230.2010.03967.x.
5. Michel F, Navellou JC, Ferraud D, Toussiroit E, Wendling D. "DRESS syndrome in a patient on sulfasalazine for rheumatoid arthritis." *Joint Bone Spine*. 2005 Jan; 72(1):82-5.
6. Kardaun SH, Sidoroff A, Valeyrie-Allanore L, et al. "Variability in the clinical pattern of cutaneous side-effects of drugs with systemic symptoms: does a DRESS syndrome really exist?" *Br J Dermatol*. 2007; 156:609-611.
7. Shiohara T, Inaoka M, Kano Y. "Drug-induced hypersensitivity syndrome (DIHS): a reaction induced by a complex interplay among herpesviruses and antiviral and antidrug immune responses." *Allergol Int*. 2006; 55:1-8.
8. Camous X, Calbo S, Picard D, Musette P. "Drug Reaction with Eosinophilia and Systemic Symptoms: an update on pathogenesis." *Curr Opin Immunol*. 2012 Dec; 24(6):730-5.
9. Picard D, Janela B, Descamps V, D'Incan M, Courville P, Jacquot S, Rogez S, Mardivirin L, Moins-Teisserenc H, Toubert A, Benichou J, Joly P, Musette P. "Drug reaction with eosinophilia and systemic symptoms (DRESS): a multiorgan antiviral T cell response." *Sci Transl Med*. 2010 Aug 25; 2(46):46ra62.
10. Fields KS, Petersen MJ, Chiao E, Tristani-Firouzi P. "Case reports: treatment of nevirapine-associated dress syndrome with intravenous immune globulin (IVIG)." *J Drugs Dermatol*. 2005 Jul-Aug; 4(4):510-3.
11. Ponnusamy Santhamoorthy, K John Alexander, and Asmahan Alshubaili. "Intravenous immunoglobulin in the treatment of drug rash eosinophilia and systemic symptoms caused by phenytoin." *Ann Indian Acad Neurol*. 2012 Oct-Dec; 15(4): 320-322.
12. Walsh S, Diaz-Cano S, Higgins E, Morris-Jones R, Bashir S, Bernal W, Creamer D. "Drug reaction with eosinophilia and systemic symptoms: is cutaneous phenotype a prognostic marker for outcome? A review of clinicopathological features of 27 cases." *Br J Dermatol*. 2013 Feb; 168(2):391-401.
13. Chen YC, Chang CY, Cho YT, Chiu HC, Chu CY. "Long-term sequelae of drug reaction with eosinophilia and systemic symptoms: A retrospective cohort study from Taiwan." *J Am Acad Dermatol*. 2013 Mar; 68(3):459-65.
14. Ushigome Y, Kano Y, Ishida T, Hirahara K, Shiohara T. "Short- and long-term outcomes of 34 patients with drug-induced hypersensitivity syndrome in a single institution." *J Am Acad Dermatol*. 2012 Nov 19. pii: S0190-9622(12)01144-9.
15. Criado PR, Avancini J, Santi CG, Medrado AT, Rodrigues CE, de Carvalho JF. "Drug reaction with eosinophilia and systemic symptoms (DRESS): a complex interaction of drugs, viruses and the immune system." *Isr Med Assoc J*. 2012 Sep;14(9):577-82.

16. Ogawa K, Morito H, Hasegawa A, Daikoku N, Miyagawa F, Okazaki A, Fukumoto T, Kobayashi N, Kasai T, Watanabe H, Sueki H, Iijima M, Tohyama M, Hashimoto K, Asada H. "Identification of thymus and activation-regulated chemokine (TARC/CCL17) as a potential marker for early indication of disease and prediction of disease activity in drug-induced hypersensitivity syndrome (DIHS)/drug rash with eosinophilia and systemic symptoms (DRESS)." *J Dermatol Sci.* 2013 Jan; 69(1):38-43.

ASK A PATHOLOGIST

Emily Coberly, MD, Krystal Tray, MT, and William Miller, MD

University of Missouri– Columbia

QUESTION: A 53 years old female receives transfusion of 2 units of PRBCs. One hour later, she develops acute dyspnea with decreased O₂ saturations. Chest x-ray demonstrates new bilateral pulmonary infiltrates. Could this patient have TRALI?

ANSWER: Yes, Transfusion Related Acute Lung Injury (TRALI) is an uncommon but extremely serious complication of blood product transfusion. It can occur with transfusion of any blood products that contains plasma, including RBCs, FFP, whole blood, platelets, and cryoprecipitate. The risk of TRALI is estimated to be between 1:2000 and 1:5000 units of blood products; and TRALI is the leading cause of transfusion-related mortality.

TRALI is most commonly caused by donor plasma containing anti-leukocyte antibodies, which react with neutrophils in the pulmonary micro-vasculature of the recipient, leading to increased vascular permeability and a pulmonary capillary leak syndrome resembling ARDS.

By definition, TRALI occurs during or within 6 hours of transfusion of a plasma-containing blood product. Symptoms include sudden onset of respiratory distress with clinical and x-ray evidence of acute bilateral pulmonary edema. Hypotension and fever are often present. TRALI is a diagnosis of exclusion and must be distinguished from the more common Transfusion Associated Circulatory Overload (TACO). BNP and NT-pro-BNP levels can be elevated in both conditions and do not reliably distinguish between the two diagnoses. As opposed to TACO, TRALI does not respond to diuretics. Treatment of TRALI is supportive; symptoms typically resolve within 72 hours. Mortality is 5-10%.

If you suspect a patient may have TRALI, it is extremely important to contact your pathologist or blood bank immediately. Since TRALI is often caused by antibodies in donor plasma, the blood donor center must be contacted to quarantine any remaining products from the suspect donors. Suspect donors are tested for anti-leukocyte antibodies, and donors implicated in an episode of TRALI are evaluated for continued eligibility to donate blood products.

Useful References:

1. Curtis BR, McFarland JG. Mechanisms of Transfusion-Related Acute Lung Injury (TRALI): Anti-Leukocyte Antibodies. *Crit Care Med.* 2006; 34(5): S118-S123.
2. Guangxi L, Daniels CE, Kojicic M, Krpata T, Wilson GA, Winters JL, Breannan Moore S, Gajic O. The Accuracy of Natriuretic Peptides (BNP and NT-pro-BNP) in the Differentiation between