

MU Guide

Swine Antibiotics and Feed Additives: Food Safety Considerations

Marcia S. Carlson and Thomas J. Fangman
Department of Animal Sciences

The term *antibiotic* means “against life” or “destructive to life.” An antibiotic is a natural compound synthesized by a living organism that inhibits the growth of another living organism.

Nearly all antibiotics are produced by bacteria or molds. Many microorganisms produce antibiotics that inhibit or kill other organisms, a process called *antibiosis*. Some antibiotics are *bacteriostatic*; they prevent growth of bacteria. Others, like penicillin, are *bactericidal*; they destroy bacteria. Groups of antibiotics based on their action are as follows:

- Agents that act on the cell wall of bacteria, such as penicillin and bacitracin.
- Agents that have a detergent effect on the cell membrane, such as polymyxin and novobiocin.
- Agents that interfere with protein synthesis, such as tetracyclines and streptomycin.
- Agents that affect nucleic acid metabolism, such as griseofulvin.

Another way to compare antibiotics is to look at their scope of effectiveness.

- Narrow-spectrum antibiotics have an antibacterial effect on a relatively small number of species.
- Broad-spectrum antibiotics are active against a variety of organisms. In general, where no specific disease is isolated, the antibiotic with the broader spectrum — that is, effective against the greater number of detrimental organisms — is preferred.

Antibiotics have a short history. In 1929, Alexander Fleming discovered penicillin by accident when a small amount of penicillin fungus landed on a petri dish loaded with staphylococci. During the 1930s, penicillin could be obtained only from government sources and was used to treat soldiers with infections or gangrene. Penicillin was first used in the U.S. general public in 1942 to combat staphylococci among victims of a nightclub fire. Over the next 50 years, scientists and health professionals learned to make use of the therapeutic and prophylactic qualities of many antibiotics as they



Proper use of feed additives can help ensure swine health and growth performance.

became available. In swine production, the development of a large number of antibiotics and other additives has helped increase gain and reduce the feed required per unit of gain.

At least 11 antibacterial or antifungal compounds or groups of compounds are widely used in swine feeds. These compounds include various salts of bacitracin, chlortetracycline, dynafac, mycostatin, oxytetracycline, oleandomycin, penicillin, streptomycin, bambermycins, tilmicosin and tylosin.

Chemotherapeutics and anthelmintics

Chemotherapeutics are organic compounds with bacteriostatic or bactericidal properties similar to those of antibiotics. But, unlike antibiotics, these compounds are produced chemically rather than microbiologically. Anthelmintics, or dewormers, are also organic compounds added to swine diets generally for short intervals to help control worms in growing-finishing swine and the breeding herd.

Other additives with potential for swine feeds are also commonly used, such as carbadox and sulfas. New products are likely to continue to be developed and approved.

Mode of action

The referring veterinarian commonly uses feed-grade antibiotics at therapeutic levels to treat an acute disease outbreak following a diagnosis. In this way, the antibiotic sensitivity of the bacterial pathogen causing the observed clinical signs is treated with an antibiotic to which the pathogen has a demonstrated susceptibility. This mode of specific pathogen treatment results in a bactericidal or bacteriostatic effect; that is, the treatment either kills the pathogen or prevents it from spreading.

In many instances, antibiotics are used to promote growth, or weight. It is generally accepted that antibiotics aid in the promotion of quality pork growth through a direct metabolic effect, by sparing use of nutrients or by controlling low levels of endemic pathogens.

1. Nutrient sparing effect

The effects of antibiotics on nutrition have been the subject of extensive research since the early 1950s. In hogs, the action of antibiotics in a nutrient sparing role is confined to the effect on the microbial population of the intestinal tract, particularly the large intestine and the lower regions of the small intestine. Microbial activity affecting nutrition in the stomach is very limited.

Antibiotics can influence nutrition in the following ways:

- Enhancing the growth of intestinal organisms that synthesize nutrients required by the animal. Such organisms may provide vitamins and amino acids. They may also digest cellulose (fiber) to end products that are useful to the animal. Pigs, like other monogastric animals, cannot digest fiber.
- Depressing growth of organisms that compete with the host animal for nutrients. Such organisms include those which break down glucose, degrade amino acids and alter fatty acids and bile salts.
- Reducing the intestinal wall thickness, implying the potential for improved absorption.

2. Disease control effect

By suppressing disease-causing organisms, including toxin producers, in the animal's environment, antibiotics may reduce the incidence of clinical and subclinical diseases that hinder performance. Producers may see these disease control effects in a greater response to antibiotics in the following situations:

- In environments that are not intensively managed for cleanliness.
- In hogs that are gaining at slower rates.
- In buildings where mixed-age swine are present.
- In older buildings.

3. Metabolic effect

The metabolic effect of an antibiotic is defined as a response that alters the rate or pattern of some biochem-

ical process at the tissue (muscle) level. Most growth-promoting antibiotics are not present in high enough concentrations at the tissue level to have a bacteriostatic or bactericidal effect on bacteria.

Choosing a feed additive

Several growth-promoting feed additives are available to swine producers. Table 1 lists several commonly used feed additives, their approved levels, and their withdrawal times.

Consider the required withdrawal time before slaughter when choosing a feed additive. Some feed additives remain in tissues longer than others. The level fed and the duration of feeding also influence tissue retention of additives. Abide by the required withdrawal times and use only the approved concentrations.

Also consider your farm's environment, management conditions, and the stage of production cycle when choosing a feed additive. These things will cause the specific feed additive and the level needed for the best response to vary.

The degree of response to feed additives will vary with the control of disease organisms in the pig's environment. Response will also vary by stage of production; in certain stages, response to feed additives is clearly seen; in others, a response is less evident.

The first few weeks of the pig's life are by far the most critical in terms of nutritional needs or health protection. Born with no protection against disease organisms, the pig enters an environment saturated with a variety of bacterial organisms.

The first milk, colostrum, of the sow will supply the pig with antibodies against the disease organisms in the surroundings, if the sow has been previously exposed to the organisms for enough time to synthesize the antibodies and concentrate them in the colostrum. However, by three weeks of age, this acquired immunity begins to disappear. The pig does not begin producing antibodies until five or six weeks of age. Thus, between three and five weeks, the young pig is most vulnerable to disease, especially to any new, infective organism that enters the pig's environment.

Also, during the early weeks of a pig's life, it is exposed to several stress conditions that render it more susceptible to diseases: castration, weaning, treating for anemia, ear notching, vaccination, climatic stresses, and exposure to internal and external parasites. Research has shown marked responses to antibiotics during these early production stages. By the time the pig reaches 40 to 50 pounds body weight, its own disease protective system — antibody formation — is functioning well, and it has adapted to environmental stresses. This is why a practical feed additive program calls for a reduction in levels as the pig develops or progresses to market weight. Fast-growing, healthy pigs may not benefit from antibiotics or chemotherapeutics during the finishing phase. However, on high health farms, improved aver-

Table 1. Feed-grade antibiotics for targeted diseases and response.

Disease		Compound		Approved treatment levels					
				Prophylactic (g/ton)	Withdrawal time (days)	Therapeutic (g/ton)	Withdrawal time (days)	Feeding duration (days)	
Bacterial diarrhea									
E. coli	Lawsonia	Salmonella	Dysentery						
x				Apramycin	150	28		14	
		x ¹		Arsanilic acid (AA)	45–90	5	90	5	-
		x ¹		AA and bacitracin MD (BMD)	45–90 and 10–30	5			
		x ¹		AA and bacitracin zinc (BZ)	45–90 and 10–50	5			-
		x ¹		AA and C or O			45–90 and >100	5	-
		x ¹		AA and P			45–90 and 100	5	-
		x		Bacitracin MD (BMD)			250 (for G/F)	-	-
				BMD for gestating sows ²			250	-	-
		x		BMD and R			250 and 22.7–34	5	-
	x	x		Carbadox (CX)			50 (>15%CP; <75lb BW)	70	-
	x	x		CX and pyrantel tartrate			50 and 96	-	-
x	x			Chlorotetracycline (C)			10 mg/lb BW	See label	14
x	x			C and BMD			10 mg/kg BW and 10–30	See label	14
				C and hygromycin B			10 mg/kg BW and 12	15	14
		x		C and R			10–50 and 181.5	5	6
x	x			C and R			10 mg/lb BW and 22.7–34	5	14
x	x	x		C and R			10 mg/lb BW and 181.5	5	6
	x	x		C + SM + P			100 + 100 + 50	15	-
	x	x		C + ST + P			100 + 100 + 50	7	-
x	x	x		C and tiamulin H fumarate (THF)			10 mg/lb BW and 35	2	-
		x		Lincomycin	40	-	100	-	21
				Neomycin (N)	(used in combinations)				
x	x			Oxytetracycline (O)			10 mg/lb BW	5	7–14
	x	x		O and R			>100 and 22.7–34	5	-
	x	x		O and N ³			50–100 and 35–140	5	-
		x		Roxarsone (R)			181.5	5	6
				R and P			22.7–34 and 100	5	-
				Sulfamethazine (SM)	(used in combinations)				
				Sulfathiazole (ST)	(used in combinations)				
		x		Tiamulin H fumarate (THF)	35	2	200 (for 14 days)	7	-
		x		Tylosin (T)			40–100	-	21
x				T			100	-	21
		x		T and pyrantel tartrate			40–100 and 96	-	-
		x		T + SM			100	15	-
		x		Virginiamycin ⁴	25	-	50–100	-	14 (at 100 g/ton)
Bacterial respiratory disease									
Pasteurella	Mycoplasma								
x				Chlorotetracycline			10 mg/lb BW	See label	14
x				C and BMD			400 and 10–30	See label	14
x				C and R			10 mg/kg BW and 22.7–34	5	14
	x			Lincomycin			200	-	21
x	x			Oxytetracycline			10 mg/lb BW	-	7–14
x				Tilmicosin			181–363	7	-
x				T + SM			100	15	-

Table 1. Feed-grade antibiotics for targeted diseases and response. (Continued)

Disease	Compound	Approved treatment levels				
		Prophylactic (g/ton)	Withdrawal time (days)	Therapeutic (g/ton)	Withdrawal time (days)	Feeding duration (days)
Bacterial reproductive disease⁵						
Leptospirosis						
x	Chlorotetracycline			400	-	14
x	Oxytetracycline			10 mg/lb body weight	-	7–14
Cervical abscesses (Septicemia)						
x	Chlorotetracycline			50–100	-	-
x	C + SM + P			100 + 100 + 50	15	-
x	C + ST + P			100 + 100 + 50	7	-

Source: 1999 Feed Additive Compendium, Minneapolis, Minn.: Miller Publishing Co.

Notes for Tables 1 and 2.

These feeding recommendations are for growing and finishing swine unless specifically noted. Feeding duration should not exceed the number of days specified, but proper withdrawal time should still be observed.

- “x” implies the specific bacteria targeted.
- “and” implies two different compounds approved for use at the same time.
- “+” implies a ready-made manufactured combination.

1. Control of swine dysentery claim with 90 g/ton inclusion level.
2. BMD for control of clostridium perfringens in suckling pigs when fed in sow ration from 14 days pre-farrowing and 21 days post-farrowing.
3. Can be fed to baby pigs, growing-finishing pigs and sows during gestation and lactation.
4. Virginiamycin should be fed at a rate of 100 g/ton for 14 days followed by 50 g/ton for treatment of swine dysentery up to 120-pound pigs and 100 g/ton for 14 days for nonbreeding stock larger than 120 pounds.
5. Should be fed to the breeding female.
6. Should be fed to prestarter and starter pigs. (Table 2)

age daily gain can be observed when low concentrations of antibiotics are fed to alter the enteric flora and prevent subclinical levels of enteric diseases that reduce nutrient absorption, such as the endemic presence of Lawsonia (ileitis) or Salmonella species.

Proper use of feed additives

Consumers are increasingly concerned about bacterial resistance and drug residues in animal tissues. Producers should use extreme caution and follow feeding directions on the label exactly. The Food and Drug Administration regulates the use of feed additives. The Animal-Plant Health Inspection Service of the U.S. Department of Agriculture and the state feed regulatory agencies are responsible for compliance with FDA regulations and are actively initiating more rigid controls for monitoring feed and pork at packing houses for residues of feed additives. Producers should abide by the FDA regulations on removal of certain additives before selling hogs for slaughter. Disregarding these regulations could cause costly losses to producers because of tissue residues and the loss of certain effective compounds for use as feed additives.

In addition to the formation required for nonmedicated feeds, the FDA requires that all medicated feeds carry the following information on the tag:

- **Purpose of the medication.** The FDA evaluates the drugs for effectiveness in growth promotion or disease prevention and treatment. Those purposes that have been adequately tested will appear on the label.
- **Directions for use.** Mix at proper levels and do not use unapproved combinations.
- **Names and amounts of all active drug ingredients.**
- **The withdrawal period.** A warning or caution for withdrawal is on the label when required for the particular drug contained in the feed.
- **Warning against misuse.** This will explain the adverse effects of using too high levels or feeding at the wrong stage of production.
If you use medicated feeds:
 - **Read the feed tag.** Be sure you’re using the medicated feed for the right stage of production and for tested and approved reasons.
 - **Meet the withdrawal times** to avoid residues and to ensure safe and wholesome pork.
 - **Do not assume withdrawal will just happen.** The time after last feeding required for a drug to clear the system varies with the type of and level of drug. All approved drugs have been tested for clearance time, and the withdrawal period is based on research.
 - **Use medicated feeds only for the purpose and**

Table 2. Feed-grade antibiotics for improvement in growth performance.

Disease/Compound	Approved treatment levels		
	Prophylactic (g/ton)	Withdrawal time (days)	Feeding duration (days)
Growth promotion–Feed efficiency (Presence of atrophic rhinitis)			
Arsanilic acid (AA)	45–90	5	-
AA and bacitracin MD (BMD)	45–90 and 10–30	5	-
AA and bacitracin zinc (BZ)	45–90 and 10–50 or 10–30	5	-
Bacitracin MD (BMD)	10–30	-	< 250 lb body weight
BMD and R	10–30 and 22.7–34	5	-
Bacitracin zinc (BZ)	10–50	-	-
Bambermycin	2–4	-	-
Carbodox	10–25	70	> 15% CP; < 75 lb body weight
Chlorotetracycline	10–50	-	-
C + SM + P	100 + 100 + 50	15	< 75 lb BW
C + ST + P ⁶	100 + 100 + 50	7	10 lb body weight up to 6 wk post-wean
Lincomycin	20	-	-
Oxytetracycline	10–50	-	-
O and N	50–150 and 70–140	5	-
Penicillin (P)	10–50	-	-
Roxarsone (R)	22.7–34	5	-
R and P	22.7–34 and 100	5	-
Tiamulin H fumarate (THF)	10	-	-
Tylosin (see production phase)			
Starter	20–100	-	-
Grower	20–40	-	-
Finisher	10–20	-	-
T and hygromycin B	10–100 and 12	15	-
T + SM	100	15	-
Virginiamycin	5–10	-	-

Source: 1999 *Feed Additive Compendium*, Minneapolis, Minn.: Miller Publishing Co.

species indicated. Drugs that are effective and approved for use in other species may not be effective in pigs, or the clearance time may differ.

- **Do not give additional drugs to animals on medicated feed without professional approval.** One drug may interfere with the effectiveness or clearance rate of another drug.
- **Do not permit other drugs to contaminate medicated or nonmedicated feed through mixer contamination or by other means.**

Although the responses to additives are more variable in gestating-lactating sows, you can expect a response during breeding and just before and after farrowing. Research has shown that feed additives in the breeding ration will increase conception rate and litter size, and additives in the farrowing ration increase pig survival and performance.

For certain diseases and parasites, feed additives can control a specific problem through short-term use of higher levels than those allowed for growth promotion. Correctly diagnosing the trouble and matching the additives to the problem are important. Table 1 contains a list of specific diseases and the feed-grade antibiotics

that have proved effective in controlling or treating the problem.

Producers should have their own feed additive programs as prescribed/defined by a veterinarian. Some may need to feed a certain antibiotic or chemotherapeutic compound during all stages of growth and development. Others may need to feed a certain additive or combinations for a certain period, then discontinue or change to another additive for additional growth periods **only** (pulsating). Each of these programs can be effective as long as you maintain a standard feeding program, using certain antibiotics or chemotherapeutics for growth promotion while reserving others for disease outbreaks. Table 2 presents some recommended levels and compounds of antibiotics for improvement in growth performance and feed efficiency.

This does not mean that you should stay with one antibiotic or chemotherapeutic year after year, but you should avoid continual switching every two or three weeks or using a number of different antibiotics within a short period. By following a standard antibiotic or chemotherapeutic program, you and your veterinarian can plan and adapt a treatment or a preventive medicine

program accordingly.

Always follow good feeding, sanitation and disease control programs, and don't expect to buy management in a bag of medicated feed.

Summary

All pork processing facilities will be expected to have a Hazard Analysis Critical Control Point (HACCP) plan in place that will ensure that pork processed in the United States is free of salmonella contamination. There will also be a greater emphasis placed on surveillance of pork carcasses for antibiotic residues when the HACCP plans are initiated. Therefore, as producers or

advisers it is imperative that you have an understanding of the reasons that a feed additive will be formulated into the ration. The table included in this guide serves as a reminder of the legally approved use of the described feed-grade antibiotics and the mandatory withdrawal periods required for each compound. It is critical that the proper feed-grade antibiotic is selected according to the history of the farm or based on recommendations of a veterinarian for the specific pathogen or condition to be treated. It is also important that the proper withdrawal time for the feed-grade antibiotic or feed additive is strictly adhered to.

