

University of Missouri Extension

G4610, Reviewed October 1993

The Bluegrasses

Jimmy C. Henning and Howell N. Wheaton
Department of Agronomy

Kentucky bluegrass, *Poa pratensis*, is the most predominant pasture grass in the northern half of Missouri. It is adapted to well-drained loams and heavier soil types of medium and better productivity. Bluegrass will be dominant in pastures only if the soil has a salt pH of 5.3 or higher and at least a medium level of phosphorus. Although the grass often becomes dormant during dry weather, it will survive severe droughts.

Kentucky bluegrass is a rhizomatous plant that produces a dense sod. Optimum temperatures for growth range from 60 to 90 degrees Fahrenheit. Serious injury will occur if the grass is subjected to continuous soil and air temperatures of 100 degrees Fahrenheit and above. Bluegrass is relatively unproductive in midsummer, but yields can be increased and sustained by favorable moisture and nitrogen fertilization.

At comparable growth stages, bluegrass contains more energy per pound of dry matter than smooth brome grass. It is also extremely palatable during periods of rapid growth. Because of its prostrate growth habit, in early clipping studies it did not compare in yields with taller growing species such as brome grass, orchard grass or reed canary grass. However, later studies with grazing animals found it to be nearly equal with these grasses in productivity.

Most bluegrass pastures in Missouri have occurred spontaneously. The carrying capacity of these fields can be increased by weed control, addition of legumes and fertilization.

Many bluegrass fields of north and central Missouri are overgrown with weeds. This generally results from overgrazing and under fertilization. These fields can be vastly improved if you eradicate weeds and practice sound grazing management. Do not graze bluegrass lower than 2 or 3 inches. Overgrazing opens the sod to weeds and reduces the vigor and growth rate of bluegrass. Overgrazing causes very poor root and rhizome development.

Controlling pasture weeds

Pasture weeds are of three general groups: **Annuals**, **biennials** and **perennials**.

Annual species such as ragweed, fleabane and sunflower originate from seed each year and die after the seed has matured in the summer or fall.

Biennials require two years to complete their life cycle. Most thistles found in Missouri pastures and wild carrot are biennials.

Perennials originate from well established root systems that survive for many years. Seeds of perennials also produce new infestations. Ironweed, goldenrod, dock, thoroughwort and vervain are examples of this most difficult group to eradicate.

Repeated mowing or clipping may gradually reduce a stand of weeds. However, mowing must be timed

properly for good results. The early bud stage, when root reserves are low and weeds are under stress, is most favorable for effective control of biennials or perennials. Weeds mowed at this time have a slow recovery rate.

To prevent seed production, annual weeds should be mowed before seed heads mature. The optimum stage for mowing annual and perennial weeds usually does not occur at the same time.

In many cases, spraying with a herbicide is the most effective way to control broadleaf weeds growing in bluegrass pastures.

The herbicide 2,4-D is effective in controlling some broadleaf weeds. Two types of 2,4-D are available. The low volatile ester form is more expensive but more effective than the amine form.

The best time to spray for weed control is when the weeds are growing rapidly and are in the vegetative stage. Apply 2,4-D (low volatile ester) at a rate of 1 acid equivalent per acre. If perennial weeds are predominant, the rate should be increased to 2 pounds per acre.

If 2,4-D amine is used, the rate should be 1 to 2 pounds per acre. Using 2,4-D amine and mowing or clipping is an excellent way to control weeds while maintaining lespedeza in a bluegrass sod.

Use extreme care when highly sensitive crops such as tobacco, tomatoes, soybeans or alfalfa are growing near the area of application. In such situations, the amine form may be safer to use. Apply at low pressure using nozzles that discharge a coarse spray. Do not apply 2,4-D if the wind is blowing toward a sensitive crop.

Bluegrass and other forages

Legumes are often established in existing bluegrass fields to improve production and strengthen the production curve during the summer months. You can expect bluegrass with a legume to produce nearly as much beef as orchardgrass, timothy or fescue with the same legume.

Birdsfoot trefoil, ladino clover, lespedeza, red clover or alfalfa may be seeded in bluegrass sods. Trefoil is probably the best suited companion crop with bluegrass because of its heavy summer production when bluegrass is semi-dormant. Trefoil-bluegrass pastures can be expected to produce twice as much as straight bluegrass fertilized with 60 pounds of nitrogen. Production with trefoil will also be much more uniform throughout the growing season than on heavily fertilized straight bluegrass pastures.

In one seven-year test, bluegrass-trefoil produced 35 pounds more beef per year than straight bluegrass fertilized with 120 pounds of nitrogen annually. The increase was mostly due to the increased carrying capacity in the summer.

To add legumes to existing bluegrass fields, till the grass sod during late fall, winter or very early spring enough to kill approximately one-half the sod. Band-seed the trefoil or other legumes using a high analysis phosphate fertilizer such as 0-45-0.

The legume seeding should be done during January or February if it is broadcast. If it is band-seeded with a drill, it may be done as late as March. Seedings made after April 1 must compete with the grass.

Fertilizing bluegrass

Pure bluegrass sods should be fertilized annually with at least a 60-20-20 fertilizer. In fields with medium to high potash levels, the potash may be omitted. If legumes are to be maintained in a stand, the nitrogen should

be omitted and a higher level of potash and phosphate used. For example, a 0-30-60 fertilizer is well suited to bluegrass-lespedeza or trefoil pastures.

Although Kentucky bluegrass usually appears spontaneously, it is occasionally seeded as a pasture grass. When bluegrass is seeded for this purpose, common or commercial seed is used. It is usually sown in a mixture with other grass and legume seeds. Usually 2 to 4 pounds per acre of bluegrass seed are added to a normal forage mixture. When seeded alone, use 10 pounds per acre of certified quality seed. Seeding in the fall before Oct. 15 is best.

Canada bluegrass, *Poa compressa*, is also found in many Kentucky bluegrass fields in Missouri. This is usually a signal that the field is lacking in lime or phosphate or both because Canada bluegrass is more tolerant of acid, low phosphate soils than Kentucky bluegrass.

Canada bluegrass has a bluer foliage than Kentucky bluegrass, matures later, is less productive and makes very slow or little recovery after it is grazed. It is an inferior species, and fields infested with it should have a soil test taken and the soil nutrient deficiencies corrected.

Re-establishing a bluegrass sod

Although sod stripped from bluegrass pastures is usually inferior to that produced under specialized management, it is sometimes used for commercial sodding purposes.

After bluegrass has been stripped from a field it is desirable to re-seed in order to re-establish the sod. Sometimes enough rhizomes (underground stems) may remain in the soil to regenerate a sod, but this is a slow process and the stand is often spotty for many years. If some rhizomes remain, 20 pounds of seed per acre may be adequate. If not, the rate should be increased to 60 pounds per acre. Use quality seed that is free of weeds.

Seeding in mid-August to mid-September is preferred. Moderate discing provides enough seedbed preparation. Pack the seedbed with a roller or cultipacker before and after seeding. Bluegrass seed is very small — in a loose seedbed it will be covered too deeply. The seed should never be covered deeper than 1/4 inch. The following season it may be pastured, but avoid trampling and overgrazing.

When sod is stripped from a field, a large amount of plant nutrients is usually removed. Use a soil test to guide fertilizer application when re-seeding bluegrass. If soil test results are not available, work 120 pounds of phosphate and 100 pounds of potash into the seedbed. Use an additional 60 pounds of phosphate and 30 pounds of potash plus 40 pounds of nitrogen as a starter. The starter may be applied in a drill at seeding time or, if seed is broadcast, disc it into the top 2 inches of soil. The following spring, topdress with an additional 60 to 80 pounds of nitrogen.

Note

Under provisions of the Missouri Nursery Law, all sod, either native or nursery, must be inspected and an inspection certificate issued before it is sold and transported. Make application for inspection and certificate to the Office of State Entomologist, State Board of Agriculture, Jefferson City, Missouri.

G4610, reviewed October 1993

Related MU Extension publications

- G4642, Establishing Birdsfoot Trefoil in Bluegrass Sod

<http://extension.missouri.edu/publications/DisplayPub.aspx?P=G4642>

- G4650, Establishing Forages

<http://extension.missouri.edu/publications/DisplayPub.aspx?P=G4650>

- G4651, Renovating Grass Sods With Legumes

<http://extension.missouri.edu/publications/DisplayPub.aspx?P=G4651>

Order publications online at <http://extension.missouri.edu/explore/shop/> or call toll-free 800-292-0969.



■ Issued in furtherance of the Cooperative Extension Work Acts of May 8 and June 30, 1914, in cooperation with the United States Department of Agriculture. Director, Cooperative Extension, University of Missouri, Columbia, MO 65211
■ an equal opportunity/ADA institution ■ 573-882-7216 ■ extension.missouri.edu