



## Q / Which drugs are best when aggressive Alzheimer's patients need medication?

### EVIDENCE-BASED ANSWER

**A** / **ATYPICAL ANTIPSYCHOTICS ARE EFFECTIVE**; so are selective serotonin reuptake inhibitors (SSRIs), and they may be safer. Atypical antipsychotics are an effective short-term (6-12 weeks) treatment for aggressive behavior in patients with Alzheimer's disease because they consistently decrease aggression scores (strength of recommendation [SOR]: **A**, multiple randomized controlled trials [RCTs]). However, evidence of drug-related deaths in patients taking these drugs mandates weighing the benefits against the risks. SSRIs may be a safer, effective alternative (SOR: **B**, limited studies).

Evidence for the efficacy of antiepileptic agents is conflicting (SOR: **C**, inconsistent patient-oriented evidence). Valproate is ineffective for treating aggression (SOR: **C**, very small RCT).

No data exist to guide long-term medication use. All available studies lasted no longer than 12 weeks.

Nonpharmacologic therapy should be the first-line treatment for aggression in patients with Alzheimer's disease. Consider drug therapy for patients who pose an imminent threat to themselves or others.

### Evidence summary

Psychotic symptoms, including aggression, in patients with dementia are a leading cause of nursing home placement and pharmacologic treatment. RCTs have demonstrated the efficacy of atypical antipsychotics in aggressive nursing home patients.

#### Risperidone significantly reduces aggression

An RCT comparing risperidone with placebo in 345 patients found that low-dose risperidone (mean 0.95 mg/d) significantly improved aggression scores (number needed to treat [NNT]=4;  $P<.001$ ). Serious adverse events included injury, cerebrovascular events, pneumonia, and accidental overdose (number needed to harm [NNH]=13).<sup>1</sup> Other RCTs also have found risperidone to be effective in reducing aggressive behavior.<sup>2,3</sup>

#### Olanzapine is effective and well tolerated

Researchers have also studied olanzapine,

another atypical antipsychotic. A 6-week RCT of 206 elderly nursing home patients with Alzheimer's disease and psychotic or behavioral symptoms found that low-dose olanzapine (5 or 10 mg/d) decreased agitation and aggression scores (olanzapine 5 mg: NNT=5; olanzapine 10 mg: NNT=6) compared with placebo. Commonly reported adverse effects included somnolence (5 mg: NNH=5; 10 mg: NNH=5) and gait disturbance (5 mg: NNH=6; 10 mg: NNH=8).<sup>4</sup> An open-label follow-up study also found low-dose olanzapine to be well tolerated and effective in decreasing agitation and aggression scores.<sup>5</sup>

#### Weigh the benefits against the risks

The US Food and Drug Administration issued a public health advisory regarding increased mortality risk after reviewing RCTs that evaluated atypical antipsychotics in patients with dementia.<sup>6</sup> A meta-analysis of

**Brice Labruzzo Mohundro, PharmD;**  
**Karla Pope, MD, MHSA;**  
**Vincent Shaw, MD**

ULM College of Pharmacy (Dr. Mohundro); Baton Rouge General Family Medicine Residency Program, Baton Rouge, La (Drs. Mohundro, Pope, and Shaw)

**Kristin Hitchcock, MSI**  
American Academy of Orthopaedic Surgeons, Rosemont, Ill



**Atypical antipsychotics are an effective short-term treatment for aggressive behavior.**

**SSRIs may be a safer, effective alternative to antipsychotics.**

15 RCTs (N=5110) that studied olanzapine, aripiprazole, risperidone, and quetiapine in patients with dementia demonstrated a small, but increased risk of death associated with their use when compared with placebo (3.5% vs 2.3%; odds ratio=1.54; 95% confidence interval [CI], 1.06-2.23;  $P=.02$ ; NNH= 83).<sup>7</sup>

A population-based (community and long-term care facilities), retrospective cohort study of atypical and conventional antipsychotics involving 27,259 matched pairs also suggested an increased risk of death. Thirty days after beginning an atypical antipsychotic medication, increased mortality was noted when compared with no antipsychotic use in both the community cohort (adjusted hazard ratio [AHR]=1.31 [95% CI, 1.02-1.70]; NNH=500) and the long-term care cohort (AHR=1.55 [95% CI, 1.15-2.07]; absolute risk difference=1.2 percentage points; NNH=83). Conventional antipsychotics were associated with higher rates of death than atypical antipsychotics (absolute risk difference=2.6 percentage points in the community group [NNH=38] and 2.2 percentage points in the long-term care groups [NNH=45]).<sup>8</sup>

### SSRIs may be an alternative

An RCT comparing citalopram and risperidone over 12 weeks in 103 patients with dementia demonstrated similar efficacy for the 2 drugs in treating agitation. Patients receiving citalopram experienced fewer adverse effects than those receiving risperidone.<sup>9</sup> The study suggests that SSRIs may be an alternative to atypical antipsychotics.

### Carbamazepine helps, valproate doesn't

Evidence regarding the use of antiepileptic medications is conflicting. One RCT of 51 patients found carbamazepine 300 mg daily to be efficacious for short-term control of agitation with good safety and tolerability. Six weeks after beginning the study, Overt Aggression Scale scores decreased 6.7 points for carbamazepine compared with 1.9 points for placebo ( $P=.008$ ). Adverse effects, including ataxia, drowsiness, postural instability, rash, weakness, and disorientation, were more common in the carbamazepine group than the

placebo group (absolute risk increase=30%; NNH=3).<sup>10</sup>

When compared with placebo, 480 mg daily of sodium valproate for 8 weeks showed no differences in controlling aggressive behavior.<sup>11</sup> In an open-label follow-up study, aggressive behavior improved from 10.52 on the Social Dysfunction and Aggression Scale to 6.31 ( $P<.001$ ), but no improvement was observed using the Clinical Global Impression Scale for aggressive behavior. Seven deaths that authors couldn't attribute to the drug occurred. Three patients experienced drowsiness. No other adverse events were noted.<sup>12</sup>

A very small, double-blind crossover RCT (N=14) evaluated 250 to 1500 mg sodium valproate daily for 6 weeks compared with placebo. A 2-week period separated the valproate and placebo regimens. Neuropsychiatric Inventory agitation and aggression scores worsened significantly with valproate (increase of 1.43 points compared with a decrease of 2.08 points with placebo;  $P=.04$ ). Adverse events related to valproate included falls, sedation, loss of appetite, thrombocytopenia, and loose stools (NNH=3).<sup>13</sup>

### Recommendations

The Expert Consensus Guideline for the Treatment of Agitation in Older Persons with Dementia<sup>14</sup> and treatment guidelines for Alzheimer's disease and other dementias from the American Psychiatric Association (APA)<sup>15</sup> offer different recommendations for first-line treatment.

The Expert Consensus Guideline recommends divalproate, risperidone, and conventional high-potency antipsychotics for patients with severe anger and physical aggression. Alternative treatments include olanzapine, carbamazepine, trazodone, and SSRIs.<sup>14</sup>

The APA recommends antipsychotics to treat agitation based on available evidence. If treatment fails, consider anticonvulsants, lithium, or beta-blockers. The APA notes that although evidence for SSRIs is limited, they may be appropriate for agitated nonpsychotic patients.<sup>15</sup>

JFP

CONTINUED ON PAGE 604

# CLINICAL INQUIRIES

CONTINUED FROM PAGE 596

## References

1. Brodaty H, Ames D, Snowdon J, et al. Risperidone for psychosis of Alzheimer's disease and mixed dementia: results of a double-blind, placebo-controlled trial. *Int J Geriatr Psychiatry*. 2005;20:1153-1157.
2. Katz IR, Jeste DV, Mintzer JE, et al. Comparison of risperidone and placebo for psychosis and behavioral disturbances associated with dementia: a randomized, double-blind trial. Risperidone Study Group. *J Clin Psychiatry*. 1999;60:107-115.
3. Frank L, Kleinman L, Ciesla G, et al. The effect of risperidone on nursing burden associated with caring for patients with dementia. *J Am Geriatr Soc*. 2004;52:1449-1455.
4. Street JS, Clark WS, Gannon KS, et al. Olanzapine treatment of psychotic and behavioral symptoms in patients with Alzheimer's disease in nursing care facilities: a double-blind, randomized, placebo-controlled trial. The HGEU Study Group. *Arch Gen Psychiatry*. 2000;57:968-976.
5. Street JS, Clark WS, Kadam DL, et al. Long-term efficacy of olanzapine in the control of psychotic and behavioral symptoms in nursing home patients with Alzheimer's dementia. *Int J Geriatr Psychiatry*. 2001;16(suppl 1):S62-S70.
6. US Food and Drug Administration. Deaths with antipsychotics in elderly patients with behavioral disturbances. Available at: [www.fda.gov/Drugs/DrugSafety/PublicHealthAdvisories/ucm053171.htm](http://www.fda.gov/Drugs/DrugSafety/PublicHealthAdvisories/ucm053171.htm). Accessed October 20, 2009.
7. Schneider LS, Dagerman KS, Insel PI. Risk of death with atypical antipsychotic drug treatment for dementia: meta-analysis of randomized placebo-controlled trials. *JAMA*. 2005;294:1934-1943.
8. Gill SS, Bronskill SE, Normand ST, et al. Antipsychotic drug use and mortality in older adults with dementia. *Ann Intern Med*. 2007;146:775-786.
9. Pollock BG, Mulsant BH, Rosen J, et al. A double-blind comparison of citalopram and risperidone for the treatment of behavioral and psychotic symptoms associated with dementia. *Am J Geriatr Psychiatry*. 2007;15:942-952.
10. Tariot PN, Erb R, Podgorski CA, et al. Efficacy and tolerability of carbamazepine for agitation and aggression in dementia. *Am J Psychiatry*. 1998;155:54-61.
11. Sival RC, Jaffmans PM, Franssen PA, et al. Sodium valproate in the treatment of aggressive behaviour in patients with dementia: a randomized, placebo-controlled clinical trial. *Int J Geriatr Psychiatry*. 2002;17:579-585.
12. Sival RC, Duivenvoorden HJ, Jansen PA, et al. Sodium valproate in aggressive behaviour in dementia: a twelve-week open label follow-up study. *Int J Geriatr Psychiatry*. 2004;19:305-312.
13. Herrmann N, Lanctot KL, Rothenburg LS, et al. A placebo-controlled trial of valproate for agitation and aggression in Alzheimer's disease. *Dement Geriatr Cogn Disord*. 2007;23:116-119.
14. Treatment of agitation in older persons with dementia. The Expert Consensus Guideline Series. *Postgrad Med*. 1998 March; Spec No:1-88.
15. Rabins PV, Blacker D, Rovner BW, et al. Treatment of patients with Alzheimer's disease and other dementias, 2nd ed. Available at: [www.psychiatryonline.com/pracGuide/pracGuideTopic\\_3.aspx](http://www.psychiatryonline.com/pracGuide/pracGuideTopic_3.aspx). Accessed October 18, 2009.

**No data exist to guide long-term medication use for aggressive behavior in patients with Alzheimer's disease.**

Visit us online at  
**[jfponline.com](http://jfponline.com)**

THE JOURNAL OF  
**FAMILY PRACTICE**

UNITED STATES POSTAL SERVICE®			Statement of Ownership, Management, and Circulation (Requester Publications Only)		
1. Publication Title The Journal of Family Practice	2. Publication Number 0 0 9 4 3 5 0 9	3. Filing Date 9/22/10			
4. Issue Frequency Monthly	5. Number of Issues Published Annually 12	6. Annual Subscription Price (if any) \$154.00			
7. Complete Mailing Address of Known Office of Publication (Not printer) (Street, city, county, state, and ZIP+4®) Quadrant HealthCom Inc., 7 Century Drive, Suite 302, Parsippany, Morris County, NJ 07054-4609			Contact Person Dorina Sckles Telephone (include area code) 973-206-6005		
8. Complete Mailing Address of Headquarters or General Business Office of Publisher (Not printer) 7 Century Drive, Suite 302, Parsippany, NJ 07054-4609					
9. Full Names and Complete Mailing Addresses of Publisher, Editor, and Managing Editor (Do not leave blank) Publisher (Name and complete mailing address) Christy Tetterton, 7 Century Drive, Suite 302, Parsippany, NJ 07054-4609 Editor (Name and complete mailing address) Marya Ostrowski, 7 Century Drive, Suite 302, Parsippany, NJ 07054-4609 Managing Editor (Name and complete mailing address)					
10. Owner (Do not leave blank. If the publication is owned by a corporation, give the name and address of the corporation immediately followed by the names and addresses of all stockholders owning or holding 1 percent or more of the total amount of stock. If not owned by a corporation, give the names and addresses of the individual owners. If owned by a partnership or other unincorporated firm, give its name and address as well as those of each individual owner. If the publication is published by a nonprofit organization, give its name and address.) Full Name Complete Mailing Address Quadrant HealthCom, Inc. 7 Century Drive, Suite 302, Parsippany, NJ 07054 Quadrant Media Corporation 450 Park Avenue, New York, NY 10022					
11. Known Bondholders, Mortgagees, and Other Security Holders Owning or Holding 1 Percent or More of Total Amount of Bonds, Mortgages, or Other Securities. If none, check box: <input type="checkbox"/> None Full Name Complete Mailing Address					
12. Tax Status (For completion by nonprofit organizations authorized to mail at nonprofit rates) (Check one) The purpose, function, and nonprofit status of this organization and the exempt status for federal income tax purposes: <input type="checkbox"/> Has Not Changed During Preceding 12 Months <input type="checkbox"/> Has Changed During Preceding 12 Months (Publisher must submit explanation of change with this statement)					
PS Form 3526-R, September 2007 (Page 1 of 3 (Instructions Page 3)) PSN: 7530-09-000-8855 PRIVACY NOTICE: See our privacy policy on <a href="http://www.usps.com">www.usps.com</a>					

13. Publication Title The Journal of Family Practice		14. Issue Date for Circulation Data Below September 2010	
15. Extent and Nature of Circulation		Average No. Copies Each Issue During Preceding 12 Months	No. Copies of Single Issue Published Nearest to Filing Date
a. Total Number of Copies (Net press run)		97,014	97,001
b. Legitimate Paid and/or Requested Distribution (By Mail and Outside the Mail)	(1) Outside County Paid/Requested Mail Subscriptions stated on PS Form 3541 (Include direct written request from recipient, telemarketing and Internet request from recipient, paid subscriptions including nominal rate subscriptions, employer requests, advertiser's proof copies, and exchange copies.)	51,299	55,641
	(2) In-County Paid/Requested Mail Subscriptions stated on PS Form 3541 (Include direct written request from recipient, telemarketing and Internet requests from recipient, paid subscriptions including nominal rate subscriptions, employer requests, advertiser's proof copies, and exchange copies.)	-	-
	(3) Sales Through Dealers and Carriers, Street Vendors, Counter Sales, and Other Paid or Requested Distribution Outside USPS®	-	-
	(4) Requested Copies Distributed by Other Mail Classes Through the USPS (e.g. First-Class Mail®)	-	-
c. Total Paid and/or Requested Circulation (Sum of 15b (1), (2), (3), and (4))		51,299	55,641
d. Non-requested Distribution (By Mail and Outside the Mail)	(1) Outside County Non-requested Copies Stated on PS Form 3541 (include Sample copies, Requests Over 3 years old, Requests induced by a Premium, Bulk Sales and Requests including Association Requests, Names obtained from Business Directories, Lists, and other sources)	45,015	40,409
	(2) In-County Non-requested Copies Stated on PS Form 3541 (include Sample copies, Requests Over 3 years old, Requests induced by a Premium, Bulk Sales and Requests including Association Requests, Names obtained from Business Directories, Lists, and other sources)	-	-
	(3) Non-requested Copies Distributed Through the USPS by Other Classes of Mail (e.g. First-Class Mail, Nonrequestor Copies mailed in excess of 10% Limit mailed at Standard Mail® or Package Services Rates)	-	-
	(4) Non-requested Copies Distributed Outside the Mail (Include Pickup Stands, Trade Shows, Showrooms and Other Sources)	-	-
e. Total Non-requested Distribution (Sum of 15d (1), (2), (3), and (4))		45,015	40,409
f. Total Distribution (Sum of 15c and e)		96,314	96,050
g. Copies not Distributed (See Instructions to Publishers #4, (page #3))		700	951
h. Total (Sum of 15f and g)		97,014	97,001
i. Percent Paid and/or Requested Circulation (15c divided by 15f times 100)		53.3%	57.9%
16. Publication of Statement of Ownership for a Requester Publication is required and will be printed in the issue of this publication. October 2010			
17. Signature and Title of Editor, Publisher, Business Manager, or Owner 			Date 9/22/10
I certify that all information furnished on this form is true and complete. I understand that anyone who furnishes false or misleading information on this form or who omits material or information requested on the form may be subject to criminal sanctions (including fines and imprisonment) and/or civil sanctions (including civil penalties).			
PS Form 3526-R, September 2007 (Page 2 of 3)			